

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Keep out of reach of children. Harmful if swallowed. Do not apply directly to food. Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After spraying in meat packaging plants, bakeries and other food processing plants, wash all equipment where exposed food will be handled with an effective cleaning compound followed by a potable water rinse. Remove pets, birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL & CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

1. Prohibitions
Do not contaminate water, food, or feed by storage or disposal.
2. Pesticide Disposal
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
3. Container Disposal
Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Space Spray — Use undiluted to kill accessible stages of insects* infesting dried fruit products in storage or being processed.

Close room or warehouse and shut off all ventilating systems. Use a spraying device such as the Micro-Gen equipment to dispense this product as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use a dosage not to exceed 1/2 gallon of this spray per 50,000 cubic feet of room space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry.

*Indian Meal Moth, Dried Fruit Beetle, Confused Flour Beetle, Saw-Toothed Grain Beetle and Roaches

CODE 16089

GCN 100202

ULD® BP-300 INSECTICIDE

For use in and around restaurants, schools, hotels, motels, apartments, commercial buildings, factories, campgrounds, resorts, and food processing establishments

ACTIVE INGREDIENTS:

Pyrethrins	3.00%
** Piperonyl butoxide, technical	6.00%
** N-octyl bicycloheptene dicarboximide	10.00%

* Consists of 4.00% (butylcarbityl) (6-propylpiperonyl) ether and 1.20% related compounds. **MGK 264 insecticide Synergist.

INERT INGREDIENTS: 81.00%
Contains Petroleum distillate.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

**MICRO-GEN
EQUIPMENT CORP.**

EPA REG. NO. 11540-1
EPA EST. NO. 11540-TX-1

Ultra Low Dosage Through Micron Generation

•• DIRECTIONS, continued

Space Spray — Use undiluted to kill the adult and accessible forms of insect pests* of (stored in multiwall paper bags or in cloth bags) in warehouses, storage rooms and similar locations: Close room or warehouse and shut off all ventilating systems. Use a spraying device such as the Micro-Gen equipment to dispense this product. Use a dosage of one fluid ounce of this spray per 1,000 cubic feet of room space. Repeat application as necessary for insect control. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry.

*Among these are the Rice Weevil, Yellow Meal Worm, Cadelle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Angourmous Grain Moth, Mediterranean Flour Moth and Indian Meal Moth

Space Spray — Use diluted to kill flies, mosquitoes and gnats in open areas:

Dilute concentrate with petroleum distillate at the rate of one part concentrate to two parts oil and mix well. Use at a time when the temperature is cool (75°F or less) and the wind velocity is approximately five MPH. With equipment such as the Micro-Gen equipment, spray at a dosage rate of fifteen ounces per minute upwind of the area to be controlled. If the equipment is truck mounted, keep the speed to approximately five MPH. Repeat application as required.

Space Spray — For control of flying and crawling insects in establishments operating under the Federal meat, poultry, shell egg grading and egg products inspection programs:

Dilute concentrate with a white mineral oil conforming to regulation 21 CFR 121.2589(b) or an odorless light petroleum hydrocarbon conforming to 121.1182, at the rate of one part concentrate to two parts oil, and mix well. Do not apply while food processing is underway. Cover or remove any food. Cover equipment that will be in direct contact with food. Using Micro-Gen equipment adjusted to deliver aerosol droplets (30 microns or less), apply one ounce per 1,000 cubic feet of space. Leave room closed for one hour and ventilate before re-entry. Before processing begins, wash down all equipment, benches, shelving etc., where exposed food will be handled, with an effective cleaning compound followed by a potable water rinse.

Space and Contact Spray — Use undiluted to kill roaches, silverfish, waterbugs, earwigs, ants, spiders, and beetles in sheltered situations: With equipment described above, apply the spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Do not allow spray to contact stored food. Use at a dosage rate of approximately one-half fluid ounce per 1,000 cubic feet of room space. Repeat dosage as necessary for control. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry.

Net Contents

Gallons

BATCH NO.

[Empty box for batch number]