

Use and Directions for Application

Lactating Dairy and Beef Animals: Flies (Horn Fly, Stable Fly, Face Fly, House Fly)

For the control of horn flies, stable flies, face flies, and house flies, make a light application daily as a fine mist to cover thoroughly all parts of the animal, including the legs.

Lactating Dairy and Beef Animals: Lice

Make 1-2 applications of 3-3½ ounces at 14 day intervals as a fine mist spray to thoroughly cover all parts of the animal. If material is applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

Back and Face Rubbers:

For the control of horn flies and face flies on lactating dairy and beef animals, use in back rubbers and face rubbers. Use 1 gallon for each 20 linear feet of burlap back rubber. For automatic back rubbers recharge the reservoir as needed. Animals must rub face on rubbers for best face fly control.

CAUTION: Harmful if swallowed.

Avoid Prolonged or Repeated inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention.

Wash thoroughly with soap and water after handling and before eating or smoking.

Do not contaminate feed, water or foodstuffs.

If swallowed, induce vomiting immediately and get medical attention.

Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Keep away from open flame.

Do not reuse container. Destroy when empty.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Product of U.S.A.

Koos, Inc. • Kenosha, Wisconsin 53140