

Cameraman's Note
Poor Copy

Net weight 20 lbs.

Ac
Lawn Food
with

Weed Control

22 - 6 - 8

Minimum Guaranteed Analysis

Total Nitrogen (N) 22%

4% Ammoniacal Nitrogen

16% Derived from Urea

Available Phosphoric Acid (P_2O_5) 6%

Derived from 54% Phosphoric Acid

Soluble Potash (K_2O) 8%

Derived from Muriate of Potash

(Potential Acidity Equivalent to 500 lbs. of calcium carbonate
per ton.)

Active Ingredients:

Dicamba

(2-methoxy-3, 6-dichlorobenzoic acid) 0.12%

2, 4-Dichlorophenoxyacetic acid 0.80%

Inert Ingredients:

Total

99.08%

100.00%

kills weeds and feeds grass

Controls Dandelions, Plantain, Buckhorn and other broadleaf weeds.
Chickweed (common, mossy and stitchwort) Clover, Curley Dock,
Dog Fennel, Knotweed, Sheep Sorrel, Red Sorrel, Creeping Charlie,
Purslane and Blackmedic.

Instructions

Application is recommended immediately after first mowing of lawn.

Mow grass and water lawn before applying.

Weeds are easiest to kill when temperature is 70° or higher.

Do not sow grass seed for at least 3 weeks after applying.

Do not spread on new lawns until after 2 mowings.

Do not apply on ivy or other ground broadleaf ground covers. Not for

use on St. Augustine grass or Dichondra - May cause injury to Bent grass and


Should additional weeds appear, other applications may be made later
in the year.

CAUTION: Avoid contact with skin, eyes and clothing.

Store away from seeds, fungicides, insecticides and other
fertilizers.

In case of contact, flush with plenty of water, for eyes get

to application avoid stuff on to clothing, hands, shoes and



(Potential Acidity Equivalent to 500 lbs. of calcium carbonate per ton.)

Active Ingredients:

Dicamba	
(2-methoxy-3, 6-dichlorobenzoic acid)	0.12%
2, 4-Dichlorophenoxyacetic acid	0.80%

Inert Ingredients:

	99.08%
Total	100.00%

kills weeds and feeds grass

Controls Dandelions, Plantain, Buckhorn and other broadleaf weeds, Chickweed (common, monseear and stitchwort) Clover, Curley Dock, Dog Fennel, Knotweed, Sheep Sorrel, Red Sorrel, Creeping Charlie, Purslane and Black medic.

Instructions

Application is recommended immediately after first mowings of lawn.

Mow grass and water lawn before applying.

Weeds are easiest to kill when temperature is 70° or higher.

Do not sow grass seed for at least 3 weeks after applying.

Do not spread on new lawns until after 2 mowings.

Do not apply on ivy or other ~~ground~~ broadleaf ground covers. Not for

use on St. Augustine grass or Dichondra - May cause injury to both grass and

Should additional weeds appear, other applications may be made later in the year.

CAUTION: Avoid contact with skin, eyes and clothing

Store away from seeds, fungicides, insecticides and other fertilizers.

In case of contact, flush with plenty of water, for eyes get medical attention.

In application avoid drift on to desirable plants, shrubs and

flowers. Avoid application on or near trees, shrubs and plants

in areas where their roots may extend and in locations where the chemical may be washed or moved.

HOLIDAY FLOWER BED WEED CONTROL

An All-Season Weed Preventer for established
Rose Gardens, Flower Beds, Shrub Areas,
Planters

KILLS UNWANTED WEED SEEDS AS THEY SPROUT

LASTS ALL SEASON

HARMLESS TO ESTABLISHED FLOWERS AND SHRUBS

DOES NOT LEACH FROM SOIL

COVERS 125 SQUARE FEET

Active Ingredients:

Trifluralin (a, a, a-trifluoro-2, 6-dinitro-N,
N-dipropyl-p-toluidine)* 1.15%

Inert Ingredients: 98.85%

100.00%

*Treflan®—The registered trademark for Trifluralin,
Elanco Products Co., a division of Eli Lilly Company.

Net Weight:
1 Lb.

4500 Thirteenth Court / Kenosha, Wisconsin 53140

DIRECTIONS

READ ENTIRE LABEL BEFORE USING

When applied to the soil, controls a wide range of weed seeds as they sprout, thus keeping planted areas weed-free for a full season. Will not harm established shrubs, plants, or flowers.

INSTRUCTIONS FOR USE

1. Remove all unwanted weeds, rake the soil smooth. Will kill sprouting weed seeds, but does not kill established grasses or weeds.
2. Punch open holes in top of can, use as a shaker and spread contents of this canister on the soil as evenly as possible over 125 square feet (approximately 10 ft. by 12 ft.). Avoid skips, because weed seeds may germinate and grow in untreated spots.
3. To improve weed control, rake material carefully into top inch of soil. Subsequent shallow cultivation will not affect weed controlling action.
4. Water the treated area lightly after application; thereafter follow your normal watering schedule.
5. Normal top dressing after application with peat moss or other mulching material will not reduce the effectiveness.

WEEDS CONTROLLED

GRASSES: Crabgrass, Foxtail, Barnyardgrass, Annual Bluegrass, Goosegrass, Stinkgrass, Sandbur.

BROADLEAVED WEEDS: Purslane, Puncturevine, Chickweed, Knotweed, Lambsquarter, Pigweed, Carelessweed, Spotted Spurge.

HORTICULTURAL CAUTIONS


1. Do not apply around Amaranthus, Celosia, Portulaca, Ajuga, Myrtle, Pachysandra, and Periwinkle.
2. Do not apply to lawn areas.
3. Do not plant flower seeds in treated areas; most annual flowers may safely be transplanted into treated areas as established bedding plants.

CAUTION: Avoid inhalation of dust or contact with skin. Avoid contamination of feeds, foodstuffs, or fishponds. Wash thoroughly after using. This product is for use only on established ornamental plants. Do not use on vegetables or other edible plants.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

KOOS, INC. • 4500 - 13th Court • Kenosha, Wis. 53140

Product of U.S.A.



ACTIVE INGREDIENTS:

MCPP

2-(2-methyl-4-chlorophenoxy) propionic acid

2,4-Dichlorophenoxyacetic acid

INERT INGREDIENTS:

Total

.94%

.94%

98.12%

100.00%

ACCEPTED

1-28-72

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND
RODENTICIDE ACT
ED UNDER NO.

2491-300

CAUTION: "Keep out of the reach of children"

(See Other Cautions on Back Panel)

NET WEIGHT 20 LBS.



LAWN FOOD

WITH

WEED CONTROL

22-6-8

SPREADER SETTINGS		
Ace (1962 and later models)		6
Ace (1960-1961 model)		2
Scott's		5½
Jackson	ACCEPTED 1-28-72 UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PEST CONTROL ED UNDER NO. 2791-300	6
Cyclone		4.5
Modern		6
Central		7½
Lawn Beauty		6¼

Made in U. S. A.

Printed in U. S. A.

MANUFACTURED FOR
ACE HARDWARE CORPORATION
 CHICAGO, ILLINOIS

BY *Kaas* inc. KENOSHA,
 WISCONSIN 53140

ACTIVE INGREDIENTS:

MCPP
2-(2-methyl-4-chlorophenoxy) propionic acid .94%
2,4-Dichlorophenoxyacetic acid .94%

INERT INGREDIENTS:

Total 98.12%
100.00%

1-28-72

2491-300


CAUTION: "Keep out of the reach of children"
(See Other Cautions on Back Panel)

NET WEIGHT 20 LBS.



LAWN FOOD
WITH
WEED CONTROL
22-6-8

EPA Reg. No. 2491-300



Covers 5,000 sq. ft. (50' x 1

NEW
ACE FORMULA
Contains...

CALCIUM
SULPHUR
IRON

Added to Combat Chlorosis (yellowing) and to supply important trace elements

EPA Reg. No. 2491-300

2491-300

Covers 5,000 sq. ft. (50' x 100')

**NEW
ACE FORMULA**
Contains...

CALCIUM

SULPHUR

IRON

Combat Chlorosis (yellowing) and to supply important trace elements to turf.

GUARANTEED ANALYSIS*

Total Nitrogen (N)	22%	Calcium (Ca)	5.00%	(CaO 7.0%)
4% Ammoniacal Nitrogen		Sulphur (S)	3.00%	(SO ₃ 7.5%)
Derived from Anhydrous Ammonia		Iron (Fe)	1.00%	(Fe ₂ O ₃ 1.5%)
18% derived from Urea		Secondary Plant Food Sources		
Available Phosphoric Acid (P ₂ O ₅)	6%	Derived from Superphosphate, Ferrous		
Derived from Triple Superphosphate		Sulphate and Sulphuric Acid.		
Soluble Potash (K ₂ O)	8%			
Derived from Muriate of Potash				
Total Available Plant Food	36%			
Chlorine, not more than	7%			

* In Michigan reads, "Minimum Guaranteed Analysis"
Potential acidity equivalent to 1100 lbs. of Calcium Carbonate per ton.

ACTIVE INGREDIENTS:

MCPP

2-(2-methyl-4-chlorophenoxy) propionic acid .94%

2, 4-Dichlorophenoxyacetic acid .94%

INERT INGREDIENTS:

98.12%

Total 100.00%

KILLS WEEDS AND FEEDS GRASS

Controls: Dandelions, Plantain, Buckhorn, Common Chickweed, Clover, Curled
Dock, Knotweed, Ground Ivy, Red Sorrel, Sticwort, and English Daisy.

INSTRUCTIONS

Application is recommended immediately after first mowings of lawn or when weeds first appear.
Mow grass and water lawn before applying.

Weeds are easiest to kill when temperature is 70° or higher.

Do not sow grass seed for at least 3 weeks after applying.

Do not spread on new lawns until after 2 mowings.

Do not apply on ivy or other broadleaf ground covers. Not for use on St. Augustine grass or
Dichondra - may cause injury to Bent grass and carpetgrass.

Should additional weeds appear, other applications may be made later in the year.

Use correct setting for your spreader as shown below, apply at a uniform rate, coverage 5,000
sq. ft. per bag.

CAUTION: Avoid contact with skin, eyes and clothing.

Store away from seeds, fungicides, insecticides and other fertilizers.

In case of contact, flush with plenty of water, for eyes secure medical attention.

In application avoid drift on to desirable plants, shrubs, and flowers. Avoid applica-
tion on or near trees, shrubs and plants in areas where their roots may extend and in
locations where the chemical may be washed or moved into contact with their roots.
Clear spreader thoroughly after application but do not clean on or near trees,
shrubs and plants.

Avoid contamination of domestic or irrigation ditches.

Keep out of the reach of children.

Do not reuse container. Destroy by burying in a safe place. Do not burn.

17

SPREADER SETTINGS	
Ace (1962 and later models)	6
Ace (1960-1961 model)	2
Scott's	5½
Jackson	6
Cyclone	4.5
Modern	6
Central	7½
Lawn Beauty	6¼

Made in U. S. A.

Printed in U. S. A.

MANUFACTURED FOR
ACE HARDWARE CORPORATION
CHICAGO, ILLINOIS

BY *Kaos* inc. KENOSHA,
WISCONSIN 53140

