



HOLIDAY

25-E

PERMISSIBLE METHOXYCHLOR

1 gallon contains 2 pounds of Technic 1 Methoxychlor

Active Ingredients

Methoxychlor, Technic 1	25.1
Xylene	9.1
Paraffinic Hydrocarbons	2.5
Inert Solvents, etc.	63.3
Total	100.0

*25.1% Methoxychlor Technic 1 is equivalent to 25.1% of 2,2-bis[4-(p-methoxyphenyl)-1,1,1-trichloroethyl]propane and other isomers and reaction products.

Keep out of the reach of children, place in warm storage (50 to 80°F) and cover it to prevent evaporation before using.

Net Content: 1.00 gal.

ACCEPTED
DEC 7 1972
U.S. DEPARTMENT OF AGRICULTURE
WASHINGTON, D.C. 20250

2491-77

CAUTION:

Keep out of the reach of children.



1972 Koss Inc.

General Directions

Always mix with water. Add a small amount of water to spray tank first, then with agitator running, add required amount of Methoxychlor Spray and fill with water. It is suitable for use in high or low volume application and in shade tree insect control can be used either in mist blower or conventional hydraulic sprayers.

For Control of Insect Carriers of Dutch Elm Disease

Dormant Treatment: Apply 1% Methoxychlor Spray (50 gallons Methoxychlor Spray with 50 gallons water) when using a mist blower or apply a 2% Methoxychlor Spray (8 gallons Methoxychlor Spray with 42 gallons water) when using a hydraulic or conventional sprayer any time after the leaves drop in the fall, up to the appearance of elm flowers or leaves, usually March in the south, or April in the north. Thoroughly spray all bark surfaces. The average 50 ft. elm will require 2 to 3 gallons of spray mixture when using a mist blower, and about 25 gallons when using a hydraulic sprayer. Spray only when temperatures are above freezing and good spraying conditions prevail.

Spring Treatment when required: Apply 6% Methoxychlor Spray (25 gallons Methoxychlor Spray with 75 gallons water) when using a mist blower or apply a 1% Methoxychlor Spray (4 gallons Methoxychlor Spray with 40 gallons water) when using a hydraulic or conventional sprayer 2-1/2 to 3 weeks after appearance of elm flowers or leaves in the spring. Thoroughly cover all leaf and bark surfaces. More spray is required per tree than in dormant period. On warm days white oil should be added for mist blower sprays.

Use for control of certain external parasites of livestock - USE FOR SHEEP, GOATS, CATTLE, HORSES, PIGS, AND POULTRY. DO NOT SPRAY DAIRY ANIMALS.

For Control of Fleas - 1 quart Methoxychlor Spray to 12 1/2 gallons of water; apply 2 quarts of spray per acre for control of hornflies; for control of Short-nosed and long-nosed Cattle Lice - apply 1/2 quart per acre for control of lice; use 1 quart Methoxychlor Spray to 12 1/2 gallons of water applied to the inside of the ear for fly control on Beef Cattle - for control of hornflies use 1 gallon Methoxychlor Spray with 12 1/2 gallons of water per acre. Use 1 gallon of solution to saturate hair on neck and face of sheep and goats every 2 to 3 weeks.

CAUTION: Barbiturate allowed. Avoid contact with eyes. Avoid contact with skin. Avoid contact with mouth. Avoid contamination of food and feed materials. Keep out of the reach of children. Do not spray or treat when temperatures exceed 85° F.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

This product is toxic to fish. Keep out of reach of streams and ponds.

Do not use in empty drums. Return to dealer for refilling or destruction of empty drums. Do not use in empty drums. Do not use in empty drums.

1) 2-15-77
2491-317