

front copy

4/14/83


GREEN TURF

WEED & FEED

20 - 4 - 6

for St. Augustine and Floratam Lawns

covers 3,000 sq. ft.



Keep out of the reach of children

CAUTION

(See back panel for additional cautions)

EPA Reg. No. 2491-

EPA Est. No. 2491-WI-1

Active Ingredient:

Atrazine (2-chloro-4 ethylamino-

6-isopropylamino-s-triazine) 0.92%

Inert Ingredients: 99.08%

Total 100.00%

Contains 18.36 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

Net Weight: 15 Lbs.

back copy: Page #2

Amount to Use: Apply at the rate of 15 lbs. per 3,000 square feet and water in well.

When to Apply: This is a preemergence material and should be applied before weeds appear in the lawn. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually applications made between January 1 and March 31, give the best control, although applications may be made through June 15, with good control of most weeds listed. To control certain winter weeds such as henbit, chickweed, rabbit tobacco, Florida betony and evening primrose, a fall application will be required.

IMPORTANT - USE PRECAUTIONS: Do Not Apply This Product on or over the Root Zone of Ornamental Trees, Shrubs, Bedding Plants, Flowers or Garden Plants or Injury to these Plants May Result.

Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result. Do not apply to newly seeded Zoysia, Centipede and Carpetgrass lawns until after the new grass has been mowed at least twice.

Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. This weed and feed may be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80°F or above, water sufficiently to rinse particles off the grass blades.

Notice: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. This weed and feed may be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80°F or above, water sufficiently to rinse particles off the grass blades.

Notice: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

Spreader Settings

Cyclone	4
Scotts	5½

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store in a clean dry place. Reseal opened bags by folding top down and securing.

Do not reuse empty container. Wrap container and place in trash collection.

KEEP OUT OF THE REACH OF CHILDREN

Manufactured by
KOOS, INC.
Kenosha, Wisconsin 53141

WEED & FEED
for
St Augustine & Floratam Lawns

Guaranteed Analysis

Total Nitrogen (N) 20%
2% Ammoniacal Nitrogen
18% Urea Nitrogen
Available Phosphoric Acid (P_2O_5) . . . 4%
Soluble Potash (K_2O) 6%

Potential acidity equivalent to 500 lbs. of
calcium carbonate per ton.

Primary Plant Food sources: Anhydrous ammonia,
Urea, Phosphoric Acid, Muriate of Potash

Chlorine, not more than 6%
431-1041

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS and DOMESTIC ANIMALS

CAUTIONS

Harmful if swallowed.
Avoid inhalation or contact with eyes. If in
eyes flush with plenty of water. Get medical
attention if irritation persists.
Do not contaminate water used for irrigation
or domestic purposes.
Do not graze treated areas.
Do not feed clippings to livestock.
Wash after handling or using.
Keep away from feed or food products.

ENVIRONMENTAL HAZARD

Keep out of lakes, streams or ponds.
Do not contaminate water by cleaning of
equipment or disposal of wastes.
Do not apply on or under ornamental trees,
shrubs, bedding plants, flowers or garden
plants or injury may result.

Harmful if swallowed.

Avoid inhalation or contact with eyes. If in eyes flush with plenty of water. Get medical attention if irritation persists.

Do not contaminate water used for irrigation or domestic purposes.

Do not graze treated areas.

Do not feed clippings to livestock.

Wash after handling or using.

Keep away from feed or food products.

ENVIRONMENTAL HAZARD

Keep out of lakes, streams or ponds.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply on or under ornamental trees, shrubs, bedding plants, flowers or garden plants or injury may result.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Weeds controlled or suppressed:

Annual Blue grass (Poa Annua)
Annual sowthistle
Annual Clover
Black Medic
Blue Toadflax
Sull Paspalum
Buckhorn plantain
Bucklover
Caroline geranium (cranesbill)
Chickweed (common & Mouseear)
Common Dandelion
Common Purslane
Crabgrass.

Cudweed
Cutleas evening Primrose
Diochondra
Dollarweed
Florida betony (rattlesnake weed)
Florida pusley
Fragrant Cudweed (rabbit tobacco)
Henbit
Knotweed
Lespedeza
Mat lippla (matchweed, creeping Charlie)
Moneywort

Narrowleaf vetch
Parsley-plant
Oxalis (common yellow wood sorrell)
Pac sorrell
Rescuegrass
Sandpur
Shephardspurse
Spurweed
Stinging nettle
Swinecrest
Venus lookingglass
Virginia pepper weed
Wild mustard