

page 1 (Oftanol 1.25%)
bag front

ACCEPTED

JUL 0 2 1987

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the product to
registered under
EPA Reg. No. 2491-330

GREEN TURF LAWN FOOD

with

Oftanol Insecticide

16-4-4

**FEEDS GRASS AND CONTROLS
LAWN INSECTS**

**CONTROLS WHITE GRUB LARVAE, CHINCH BUGS,
BILL BUGS AND MOLE CRICKETS**

(column 1)

ACTIVE INGREDIENT:

1-Methylethyl 2[[ethoxy[(1-methylethyl) amino] phosphinothioyl] oxy] benzoate].....	1.25%
INERT INGREDIENTS.....	98.75%
TOTAL.....	100.00%

(column 2)

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

**See Back Panel for Statements of Practical Treatment
and Other Precautionary Statements**

E.P.A. Reg. No. 2491-GGN

E.P.A. Est. No. 2491-WI-1

**TREATS 4,000 SQ. FT.
NET WEIGHT 15 LB. (6.8 kg.)**

***Oftanol is a Reg. TM of the Parent Company of Farbenfabriken
Bayer GmbH, Leverkusen**

Page 2 (Bag back-Oftanol 1.25%)

GREEN TURF
LAWN FOOD
with Oftanol Insecticide
16-4-4

Guaranteed Analysis

Total Nitrogen (N).....16.0%
 1.2% Ammoniacal Nitrogen
 14.8% Urea Nitrogen
Available Phosphoric Acid (P O).....4.0%
Soluble Potash (K O).....4.0%
 Primary Plant Food Sources: Ammoniated Phosphate, Urea,
 Muriate of Potash, Ammoniated Phosphates.

Chlorine (Cl), maximum.....3.8%

431-

IMPORTANT: Read these entire directions of Use and Conditions of Sale before using Green Turf Lawn Food with Oftanol Insecticide.

Green Turf Lawn Food with Oftanol Insecticide is recommended for use on lawns; and other ornamental and recreational turf grass areas.

PRECAUTIONARY STATEMENTS
Hazardous to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through the skin. May cause eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children. **SYMPTOMS OF POISONING:** A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea. In case of poisoning, call a physician or Poison Control Center. Have patient lie down and keep quiet.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give

page 3 (Oftanol 1.25%)

anything by mouth to an unconscious person. Get medical attention immediately. If on skin, wash skin immediately with soap and warm water. If irritation occurs, get medical attention. If eyes are contaminated, wash with plenty of water for at least 15 minutes. Get medical attention. If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

TO PHYSICIAN: ANTIDOTE-Atropine sulfate has a slight antidotal effect and should be administered in therapeutic doses. Repeat as necessary to the point of tolerance. Active ingredient inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RECOMMENDED APPLICATIONS

The timing of the applications vary dependent upon the insect which is being controlled. Follow the directions as specified below for control of the listed turf grass pests. Consult your local turf, State Agricultural Experiment Station, or State Extension Specialists for more specific information regarding the timing of applications.

WHITE GRUB LARVAE (such as Japanese Beetle, Black Turf Grass Ataenius, Northern masked chafer, European chafer, June and May Beetle)

Preventive Applications

To control the above listed pests apply Green Turf Lawn Food with at the rate of $3 \frac{2}{3}$ lbs./1000 sq. ft. in the Spring (April thru mid May) and immediately apply sufficient water to wet the soil 1 to 1 1/2 inches deep without causing run off.

Curative Applications

If visible damage or high grub populations have been observed apply $3 \frac{2}{3}$ lbs/1000 sq. ft. and immediately apply sufficient water to wet the soil 1 to 1 1/2 inches deep without causing run off.

page 1 (Oftanol 1.25%)

Billbug Larvae Apply $3 \frac{2}{3}$ lbs/1000 sq. ft. when adult activity is observed in the Spring usually late in March to mid May. Water in as for preventive white grub control.

Hyperion Weevil Larvae Apply at the rate of $3 \frac{2}{3}$ lbs/1000 sq. ft. in the Spring between full bloom periods for forsythia and flowering dogwood, usually April 15 to May 15 and water-in as for preventive white grub control.

Mole Crickets Apply at the rate of $3 \frac{2}{3}$ lbs/1000 sq. ft. preventively or when damage is observed. Apply 1/2 to 1 inch of water immediately after application if rainfall does not occur.

Sod Webworm Larvae Chinch Bugs Apply 1 lb. 14 ozs. to $3 \frac{2}{3}$ lbs. per 1000 sq. ft., (use higher rate for severe infestations) when first damage is first observed. Apply 1/4 to 1/2 inch of water immediately after treatment. Re-apply immediately after treatment. Re-apply as necessary for pest re-infestation (see restrictions)

RESTRICTIONS

Do not treat turf before rainfall or irrigation has been applied as per recommended applications. Keep children and pets off treated areas until dry. Do not use clippings from treated areas for feed or forage or allow food animals to graze treated areas.

The maximum amount of Green Turf Lawn Food with Oftanol Insecticide which can be used is $7 \frac{1}{3}$ lbs. per 1000 sq. ft. per year. Applications at the rate of 1 lb. 14 ozs. per 1000 sq. ft. must have a 15 day interval and applications at the rate of $3 \frac{2}{3}$ lbs per 1000 sq. ft. must have a 30 day interval between applications. A maximum 1.45 ounces of active ingredient per 1000 sq. ft. per acre per year can be applied regardless of formulation.

APPLICATION

Apply Green Turf Lawn Food with Oftanol Insecticide uniformly over the lawn area with drop-type or rotary-type spreader. Do not use a spreader that applies the product in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. The full recommended rate of Green Turf Lawn Food with Oftanol Insecticide must be used for maximum benefit.

APPROXIMATE SPREADER SETTINGS

Approximate spreader settings for applying Green Turf Lawn Food with Oftanol Insecticide at the rate of $3 \frac{2}{3}$ lbs./1000 sq. ft.

BEST AVAILABLE COPY

page 5 (Oftanol 1.25%)

MODEL

SETTING

Cyclone Model B-1
Ortho Broadcast
Ortho Whirly Bird
Scotts PF-1/011B
Scotts Broadcast

3
3
2
5
E

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PRODUCT DISPOSAL: Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Do not walk through spilled material. If material cannot be used as directed, dispose of by wrapping in paper and put in trash collection.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.

CONDITIONS OF SALE: FOLLOW DIRECTIONS CAREFULLY. THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED, OTHER CROPS, ANIMALS, MAN OR THE ENVIRONMENT. BUYER ASSUMES ALL RISKS FOR USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HERewith.

NET WEIGHT 15 LBS.

UPC

Manufactured By
KENOSHA,

0

KOON/INC.

WIS. 53140

73841