


page 1 (Oftanol 1.50%)
bag front



GREEN TURF[®]

1.5%

OFTANOL[®]
INSECTICIDE

FOR CONTROL OF LAWN INSECTS

CONTROLS WHITE GRUB LARVAE, CHINCH BUGS
BILL BUGS AND MOLE CRICKETS

(column 1)

ACTIVE INGREDIENT:

1-Methylethyl 2[[ethoxy[(1-methylethyl)
amino] phosphinothioyl] oxy] benzoate]*.....1.5%
INERT INGREDIENTS.....98.5%
TOTAL.....100.0%

(column 2)

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

**See Back Panel for Statement of Practical
Treatment and other Precautionary Statements**

EPA Reg. No. 2491-GGR

EPA EST. No. 2491-WI-1

Treats 5,000 Sq. Ft.

NET WEIGHT 15 LBS. (6.8 kg.)

*Oftanol is a Reg. TM of the Parent Company of Farbenfabriken
Bayer GmbH, Leverkusen.



2/5
page 2 (Green Turf 1.5% Oftanol)
bag back panel

GREEN TURF®

1.5% Oftanol® Insecticide

IMPORTANT: Read these entire Directions of Use and Conditions of Sale before using Green Turf® 1.5% Oftanol® Insecticide.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
If on skin: Wash skin immediately with soap and warm water.
If in eyes: flush with plenty of water for at least 15 minutes.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RECOMMENDED USES OF 1.5% OETANOL®

Green Turf® Oftanol® 1.5% Insecticide is recommended for use on lawns, and other ornamental and recreational turf grass areas.

White Grub and Billbug Larvae Control

Do not apply more than 6 lbs. of 1.5% Oftanol® per 1000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1000 sq. ft. regardless of formulation used. Minimum interval between

applications must be 30 days.

Water turf area thoroughly immediately after application, using sufficient water to wet soil 1-1/2 inches deep.

White Grub Larvae include the larvae of Japanese beetles, black turf grass atenioides, Chafers, Phyllophaga spp, Green June beetle, Oriental Beetle and Asiatic garden beetle.

Hyperodes Weevil Control (Larvae Stage):

Apply between full bloom periods of forsythia and flowering dogwood, usually April 15th to May 15th, and water-in as for Grub Larvae control.

Mole Cricket Control:

Water immediately after application using 1/2 to 1 inch of water.

Soil Webworm Larvae and Chinch Bug Control:

Apply 1/2 to 1 inch of water immediately after treatment. Reapply as necessary for pest re-infestations. Do not apply more than 6 lbs. per 1000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1000 sq. ft. regardless of formulation used. Minimum interval between applications is 15 days for 1.5 lb. rate and 30 days for the 3 lb. rate.

Area Fire Ant Control:

Apply specified dosage uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present, use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

Established Fire Ant Mounds--Drench Treatment:

Wet Conditions: Distribute dosage evenly over 9 sq. ft. and immediately sprinkle gently with 3 gallons of water.

Dry Conditions: Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment.

Make only one mound drench treatment.

NOTE: Do not apply more than 6 lbs. of 1.5% Oftanol[®] per 1,000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1,000 sq. ft. regardless of formulation used. Minimum interval between application is 15 days for 1.5 lb. rate and 30 days for the 3 lb. rate.

Consult your local turf, state Agricultural Experiment Station, or state Extension Service Specialists for more specific information regarding timing of application.

RESTRICTIONS

Do not mow treated turf area before irrigation has been applied as per RECOMMENDED APPLICATIONS. Keep children and pets off treated area until dry. Do not graze treated area

4/5
or use clippings from treated areas for feed or forage.

APPLICATION

Apply 1.5% Oftanol[®] uniformly over the lawn area with drop-type or rotary-type spreader. Do not use a spreader that applies the product in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. The full recommended rate of 1.5% Oftanol[®] must be used for maximum benefit.

APPROXIMATE SPREADER SETTINGS

Approximate spreader settings for applying 1.5% Oftanol[®] at the rate of 3 lb./1,000 sq. ft.

MODEL	SETTING
Cyclone Model B-1	3
Ortho Broadcast	3
Ortho Whirly Bird	2
Scotts PF-1/011B	5
Scotts Broadcast	E

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PRODUCT DISPOSAL: Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Do not walk through spilled material. If material cannot be used as directed, dispose of by wrapping in paper and put in trash collection.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.

CONDITIONS OF SALE: Follow directions carefully. The directions on this label were determined through research to be the directions for correct use of this product. This product has been tested for a range of weather conditions similar to those weather conditions that are ordinary and customary in the geographic area where the product is used. Insufficient control of pests may result from the occurrence of extraordinary or unusual weather, or from failure to follow label directions. In addition, failure to follow label directions may cause injury to the crop to which the product is applied, other crops, animals, man or the environment. Buyer assumes all risks for use, storage or handling of this material not in strict accordance with directions given herewith.

NET WEIGHT 15 LBS.

Manufactured by
Kenosha
KOOS/INC.
WI 53140

