


page 1 (Green Turf/Triamine Weed & Feed) front panel
K008 LABEL



PREMIUM

GREEN

TURF®

LAWN
FOOD

Provides fast and long feeding action

25-3-3

with

WEED CONTROL

Covers 5,000 Sq. Ft.

(column 1)

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichloro- phenoxyacetic Acid*.....	0.313%
Dimethylamine Salt of 2-(2-Methyl-4- Chlorophenoxy)propionic Acid**.....	0.314%
Dimethylamine Salt of 2-(2,4-Dichloro- phenoxy)propionic Acid***.....	0.310%
Inert Ingredients.....	99.063%
Total	100.000%

BY ISOMER SPECIFIC ASSAY AOAC METHOD EQUIVALENT TO:

- *0.26% of 2,4-Dichlorophenoxyacetic Acid
- **0.26% of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid
- ***0.26% of 2-(2,4-Dichlorophenoxy)propionic Acid

TRIAMINETM

Trademark of Riverdale Chemical Co.

(column 2)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Precautionary Statements

EPA Reg. No. 2491-

EPA Est. No. 2491-WI-1

NET WEIGHT 18 LBS. (8.2 kg.)

page 2 (Green Turf/Triamine)
back panel

PREMIUM

**GREEN
TURF®
LAWN FOOD
with
WEED CONTROL
25-3-3**

GREEN TURF® LAWN FOOD WITH WEED CONTROL is a stable mixture of three phenoxy herbicides. It is for use on Lawns, Ornamental Turf (not pastures) and Golf Fairways, Aprons, Tees and Roughs.

GUARANTEED ANALYSIS

431-2201

Total Nitrogen (N).....	25.00%
2.00% Ammoniacal Nitrogen	
18.80% Urea Nitrogen	
1.26% Other Water Soluble Nitrogen*	
2.94% Water Insoluble Nitrogen	
Available Phosphoric Acid (P_2O_5).....	3.00%
Soluble Potash (K_2O).....	3.00%

Primary Plant Food Sources: Anhydrous Ammonia, Urea, Methylene Urea, Phosphoric Acid, Muriate of Potash.

*1.26% Slowly Available Nitrogen from Methylenediurea and Dimethylenetriurea.

Chlorine (Cl), maximum..... 3%

This product is an effective lawn food which combines the advantage of both fast and long lasting feeding action, plus broad range weed control. The fertilizer produces a thick, green turf while killing weeds to make way for new turf growth. Weeds contacted by this product stop growing, gradually die, and disappear from the lawn.

The product is designed for control of broadleaf weeds in lawns such as:

Black Medic, Buckhorn, Burdock, Chicory, Chickweed, Clover, Dock, Dandelion, English Daisy, Ground Ivy, Heal-All, Lambsquarter, Mallow, Morning Glory, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Ragweed, Red Clover, Red Sorrel, Stitchwort, Speedwell, Spurge, White Clover, Wild Carrot, Wild Garlic, Wild Onion, Wild Lettuce, Yarrow, and other broadleaf weeds.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

page 3 (Green Turf/Triamine)

Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Causes moderate eye injury.

IF SWALLOWED: Get Medical Attention
IF ON SKIN: Wash Skin with Soap and Water
IF IN EYES: Flush Eyes with Water for 15 Minutes
and Get Medical Attention

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not apply when conditions favor drift from target areas. Drift of dust or vapor may cause injury to nearby susceptible plants such as vegetables, flowers, fruit trees and ornamentals. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DIRECTIONS

Read the entire label before using this product.

This product may be used on Blue Grass, Fescue, Rye, Bermuda, Bahia, Zoysia Lawns, Centipede and St. Augustine grass. DO NOT USE on Dichondra, and Carpet Grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in spring or early fall. IMPORTANT: Apply when grass is moist (after a rain, sprinkling or dew). DO NOT water lawn after material is applied. For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Avoid contact with valuable shrubs plants or flowers. Possible retreatment may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not overapply. Do not reseed for at least 3 to 4 weeks

Page 4 (Green Turf/Triamine)

after application. Do not apply to newly seeded or sodded lawns until after 3 mowings.

NOTE: When applying to Southern Grasses - Under conditions of adequate moisture, high temperature and high humidity, over application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 18 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader either as one pass application or a two pass application, criss-crossing the area at half the rate each time. Lawns that are heavily overrun by weeds and that are in poor condition, may require more than one application of this product. A suggested second application should be made in approximately 30 days.
DO NOT HAND SPREAD

The following settings are approximate for applying contents of this package at a suggested rate of 3.6 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. ✓

Check your spreader on 250 sq. ft. and adjust to apply at the rate of .9 lbs per 250 sq. ft.

SPREADER SETTINGS

SPREADER	ONE PASS APPLICATION		
	RATE SETTING		
Cyclone	4 3/4 - 5 1/4		
Scotts C-1	K	-	L
Scotts 35	7	-	8
Central	7		
Precision	6		

Above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Age and condition of spreader will affect application. Adjust your spreader if necessary. Be sure to shut off spreader while turning or stopping to prevent accidental over-application of grass. Fill spreader on driveway or sidewalk to eliminate chance of spillage on grass. Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

page 5 (Green Turf/Triamine)

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash. ✓

Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions provided, with proper equipment in good working order, and under normal conditions of use. If not used as directed, seller makes no warranty, expressed or implied, concerning the use of this product. User then assumes all risk of loss or damage arising from any contrary use or handling.

STORE IN A DRY PLACE

Manufactured by: Koos/Inc. - Kenosha, WI 53140

UPC

0

73841 16237