# GREEN TURF Lawn Food with Insect Control TREATS UP TO 10,000 SQ.FT.

This Product Controls larvae if the European Chafer, Japanese Beetle larvae, Hyperodes (Turfgrass) Weevil, Chinch Bugs, Sod Webworms, and other Lawns Pests, Plus Fertilizes your lawn.

See other side of bag for a complete list of insects controlled.

Treats up to 10,000 Square Feet (Read Directions for Specific Application Rates.)

ACTIVE INGREDIENT:

Chlorpyrifos: [O,O,-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate].-

INERT INGREDIENTS: - - - - - - - - - - - 99.3%

Total 100.0%

### **NET WEIGHT 33 POUNDS**

EPA Reg. No. 2491-337

# CAUTION

STATEMENTS OF PRACTICAL TREATMENT

EPA Est. No. 2491-WI-1

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomitin or give anything by mouth to an unconscious person. IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with plenty of soap an water. IF IN EYES: Flush eyes with plenty of water. NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injectio is an antidote.

#### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with eyes, skin and clothing. If watering in is directed, children and pets should be kept off treated areas until this material is washed into the soil and the grass is dry. Wash thoroughly after handling and before smoking or eating. Keep away from food, feed stuffs, and domestic water supplied.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean highwater mark. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Do not feed treated grass clippings to livestock.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your cooperative <u>Agricultural Extension service.</u>

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. <u>STORAGE</u>: Store product in its original labeled container in a cool, dry, locked place out of reach of children. <u>PESTICIDE DISPOSAL</u>: Wastes iresulting from the use of this product may be disposed of on site or at an approved waste disposal facility. <u>CONTAINER DISPOSAL</u> Completely empty bag into application equipment. Then dispose of empty lag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke <u>HOUSEHOLD</u>: Do not re use empty container. Wrap container and put in trash.

# CONTROL THESE INSECTS CUTWORMS JAPANESE BEETLES

ANTS
BROWN DOG TICKS
CHINCH BUGS

CRICKETS

EARWIGS
EUROPEAN CHAF

JAPANESE BEETLES SOD V/EBWORMS SOV/ BUGS (PILLBUGS)

HYPERODES WEEVILS

NOTE: Mosquitoes coming to rest on areas treated for control of turf pests will be killed for varying periods of time after treatment, depending on exposure to treated areas to weather conditions

### APPLICATION DIRECTIONS

Treat when insects are active or as recommended by state Agricultural Experiment Stations or State Extension Service Specialist. Apply this product uniformly to home lawns and other ornamental and recreational turf areas using any hand or power operated gravity or rotary spreader designed for application of granular materials. This product is not for pastures use. To obtain uniform distribution a apply one-half the required amount of product in one direction, then apply the balance at a right angle.

# SURFACE INSECTS

To control Ants, Brown Dog Ticks, Chinch Bugs, Crickets, Cutworms, Earwigs, Sod Webworms, and Sow Bugs, use the product at a rate of 3.3 lbs. per 1,000 sq.ft. This package will treat 10.000 sq.ft.(100' x 100') at the above rate. Do not water granules in when treating for surface insects.

## SOIL INSECTS

To control certain white grubs ( larvae of European Chafer, and Japanese Beetles,), use this product at a rate of 6 6 lbs. per 1,000 sq.ft. For best results treat when grubs are young and actively feeding near the soil surface, usually during later mid-July and August. Consult your local Agricultural Extension Service Specialist for proper time to treat in your area. Apply the granules uniformly over the area treated, followed by thorough watering to wash the insecticide into the soil.

To control Hyperodes (Turfgrass) Weevil use this product at a rate of 6.6 lbs., per 1,000 sq.ft. Treat suspected problem areas in mid April and Again in mid-May. Consult your local Agricultural Extension Service Specialist for best time to treat in your area.

## SPREADER SETTINGS

Due to variations in individual machines, these settings are intended to be a guide. It is important to make certain that your machine is spreading the correct amount of material at the setting use. The following spreader settings are basen or half rate for a two directional applications.

Make of Spreader 2 0 flor Surface Insects 5.0 Soil Insects 3.3 lbs./1,000 sq.ft 6.6 lbs./1,000 Sq.Ft. 1.5\*
Cyclone B-1 Pro 2.0\* 1.5\*
Scotts Conventional Federal Insects 6.6 lbs./1,000 Sq.Ft. 1.5\*
Scotts Conventional Insects 6.6 lbs./1,000 Sq.Ft. 1.5\*

Selier warrante the product conforms to its chemical description and it reasonably fir for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions reasonably foreseeable to seller, and buyer assumes the risk of any such use (337-amd1.pm5 \ 03-01-94)

GUARANTEED ANLYSIS

Total Nitrogen (N)→→→→ - - 1( Urea Nitrogen 10%

Available Phosphoric Acid (P205) 6%
Soluble Potash (K20) - - - - 4%

Primary plant food derived from Diammenium Phosphate, Formalderiyde Urea, Muriate of Potash

BEST AVAILABLE COPY

Kenosha, Wi 53140 U.S.A.