

DIRECTIONS

KNOX-EM SUR-DETH RAT-KIL-R is a complete rodenticide, as received. Nothing need be added.

RATS: An uninterrupted supply of bait should be maintained for approximately 10 days. Place 4 ounces to one pound of bait per bait location in feeding stations located along rat runways, walls, feeding areas, and in corners and concealed locations where rats can feed unobserved. Inspect bait everyday for the first week and at regular intervals thereafter. Replenish baits as they are eaten. Spoiled or contaminated baits should be replaced promptly. Bait should be placed in locations not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes. Baiting should continue until all signs of feeding have stopped. When reinfestation is likely to occur or a continuous source of infestation is present (nearby dumps) permanent bait stations should be established and replenished as needed.

MICE: Follow same directions for rats except that tablespoon amounts of bait should be placed at 8 to 12 ft. intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

KNOX-EM SUR-DETH RAT-KIL-R

A Ready-to-Use Rodenticide
Containing
WARFARIN

INGREDIENTS

Active Warfarin*..... 0.025%
Inert..... 99.975%
*3-(a-acetylbenzyl)-4-hydroxycoumarin

NET CONTENTS POUNDS

CAUTION

KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Cautions

THE GENERAL PEST CONTROL CO.
3561 W. 105th St. Cleveland, O. 44111
Phone: Clearwater 2-7140

USDA REGISTRATION No. 2496-4

PROPERTIES

KNOX-EM SUR-DETH RAT-KIL-R contains a powerful blood anticoagulant which is readily accepted by rodents at effective dosage levels. Highly successful rodent control has resulted when KNOX-EM SUR-DETH RAT-KIL-R has been used according to directions. Death results from internal hemorrhages if small quantities of KNOX-EM SUR-DETH RAT-KIL-R are taken over several consecutive days. Single isolated doses will not give the desired results, and therefore, larger quantities of KNOX-EM SUR-DETH RAT-KIL-R are required than when more toxic poisons, such as arsenic, thallium and strychnine are used.

PRECAUTIONS

Follow directions carefully. Place KNOX-EM SUR-DETH RAT-KIL-R in bait boxes so it will not be exposed to humans, pets, poultry or livestock. In case baits are eaten, call physician at once.

NOTE FOR PHYSICIAN: When a human is known to have ingested KNOX-EM SUR-DETH RAT-KIL-R, blood transfusions along with oral and intravenous injections of Vitamin K are indicated as in the case of hemorrhages due to overdoses of Dicumarol.

