

[2517-3]

Geisler
FOR THE LOVE OF YOUR PET

NO MITE

FOR THE CONTROL OF LICE AND MITES ON PARAKEETS AND ALL OTHER BIRDS

DIRECTIONS

1. SHAKE WELL.
2. Hold can upright at least 3 feet away from bird, press button firmly and spray for 2 seconds, directly into cage. For light infestation, spray birds twice weekly.
3. For mite infestation, remove bird from cage. Thoroughly spray perches and cage. Clean and spray at regular intervals.
4. Do not use "No Mite Spray" on plastic parts of cage or any other plastic material. May cause cloudiness.

ACTIVE INGREDIENTS:

| | |
|--|--------|
| Pyrethrins | 0.06 % |
| Technical Piperonyl Butoxide* | 0.12 % |
| N Octyl Bicycloheptene Dicarboximide** | 0.20 % |
| Petroleum Distillate | 0.29 % |
| Triethylene Glycol | 0.075% |
| Propylene Glycol | 0.075% |

.82 %

INERT INGREDIENTS:

99.18 %

100.000%

*Equivalent to 0.096% (butylcarbityl) (6-propylpiperonyl) ether and 0.024 related compounds.

** U. S. Patent No. 2,476,512.

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting. Never throw container into fire or incinerator. Cover fish and turtle bowls when spraying.



EPA REG. NO. 2517-3
EPA. EST. NO. 11715-TN-1

Item 4170

Geisler PET PRODUCTS, INC.
a *Wauqua* company

Omaha, Ne. 68105 / Bronx, N.Y. 10457

ACCEPTED
Feb. 21, 1975
E-1197-3



Geisler
FOR THE LOVE OF YOUR PET

NO MITE
SPRAY

KILLS LICE
AND MITES

NET WT. 7 OZ.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

See other cautions on back panel.

[2017-3]

Geisler

FORMULATION FOR USE ON BIRDS

Apply directly to the feathers, from head to tail, and spray for 2 seconds directly into the flight infestation. Spray birds twice weekly until fastidious remove bird from cage. This high concentration spray is not for use on mammals.

PREPARATION

1. Mix 1 part of the concentrate with 9 parts of water.

CAUTIONS

1. Do not use on mammals.

2. Do not use on birds with open wounds.

3. Do not use on birds with respiratory distress.

4. Do not use on birds with eye irritation.

5. Do not use on birds with skin irritation.

6. Do not use on birds with feather loss.

7. Do not use on birds with feather plucking.

8. Do not use on birds with feather mites.

9. Do not use on birds with feather lice.

10. Do not use on birds with feather fungus.

11. Do not use on birds with feather parasites.

12. Do not use on birds with feather damage.

13. Do not use on birds with feather loss.

14. Do not use on birds with feather plucking.

15. Do not use on birds with feather mites.

16. Do not use on birds with feather lice.

17. Do not use on birds with feather fungus.

18. Do not use on birds with feather parasites.

19. Do not use on birds with feather damage.

20. Do not use on birds with feather loss.



Geisler PET MEDICINES, INC.

Geisler

FORMULATION FOR USE ON BIRDS



FORMULATION FOR USE ON BIRDS

REACH OF WHERE