



THE LATEST EDITION OF THIS BOOK IS BASED UPON OVER 32 YEARS OF EXPERIENCE IN HANDLING AND FEEDING PETS. FOR YOUR COPY, SEND \$6.95 TO CONAGRA, 3802 LEANWORTH, DALLAS, TX 75246.



NEW
For Large Dogs

ACTIVE INGREDIENT
0,0-dimethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 15.0%
INERT INGREDIENTS 85.0%
TOTAL 100.0%

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN
See back panel for additional precautionary statements

CONTENTS: 1 FLEA & TICK COLLAR
NET WT. 0.9 OZ. (25.5g)



for Large Dogs

Adjustable - fits Large Dogs, 28" neck or smaller

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor.

Note to Physicians: Atropine is antidotal. Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs.

DIRECTIONS FOR USE

Place Flea & Tick Collar around dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of the excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn up to 150 days. Ticks are relatively harder to kill and are killed more slowly. Activity against ticks will occur within a week and collar will continue to aid in tick control in the head and neck area in unusually large breeds for up to two months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar's effectiveness is not diminished by normal wetting such as rainfall; however it is suggested collar be removed before bathing.

ACTIVE INGREDIENT
D.O. diethyl 0 (2-isopropyl 6-methyl
4-pyrimidinyl) phosphorothioate

INERT INGREDIENTS

15.0%
85.0%
100.0%

DISPOSAL: Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

GUARANTEE: If not satisfied, send used collar and indication of purchase price to ConAgra, P.O. Box 55033, Omaha, NE 68105, for refund.

Patent Pending



Manufactured in France for
ConAgra Pet Products Company
Omaha, NE 68105
Litho U.S.A.

EPA Reg. No. 257-101
EPA Est. 1557-0119
PM 4-79



0070
21953

123 12
123 123
123 123

用 器



2