

Sergeant's Dual Action Flea & Tick Collar

(with d-phenothrin)

for [Dogs], [Large Dogs], [Small Dog & Puppies], [Cats]

- [Fast Acting][Quick Kill]

- Kills [Fleas & Ticks] for 6 months

- [Deodorizes] [Deodorizing]

[Long Lasting][Gentle][Fresh][Powdery][Clean]Scent

Kills the Ticks That Carry

[Lyme Disease] [Rocky mountain Spotted Fever] [Ehrlichiosis]

[the following claims on initial labeling only; or 6 months after initial sale]

"new - improved"

"last an additional month!"

Active Ingredients:

O-Isopropoxyphenyl methylcarbamate.....9.00%

*3-Phenoxybenzyl d-cis and trans** 2,2 dimethyl-3-

(2-methylpropenyl)cyclopropane carboxylate.....3.00%

N-Octyl bicycloheptene dicarboximide.....3.00%

*d-(cis, trans) phenothrin

** cis/trans isomer ratio: max. 25% (+ or -) cis
min. 75% (+ or -) trans

Inert Ingredients:.....85.00%

Do Not Allow Children To Play With Collar

CAUTION

See cautions on back.

88: 2d 6- 100 96.

RECD LHA/GPP/DPD1

Sergeant's [Dogs] [e Dogs] [Small Dogs & Puppies][Cats]Collar

IMPORTANT CONSUMER INFORMATION

This Collar kills fleas and ticks for 6 months. Collar is [double layer] sustained-release system [containing three active agents] combining the insecticidal actions of vapors, and powders, working in [two] [three] ways to provide fast acting [quick kill] and long lasting head-to-tail protection. Thus, Collar has the effect of simultaneous usage of a spray and a powder in the control of fleas and ticks, including ticks that may carry [Rocky Mountain Spotted Fever] [Lyme Disease] [Ehrlichiosis]. Wetting Collar will not impair Collar's effectiveness. It is not necessary to remove Collar if pets go swimming or are out in the rain. Use Sergeant's Dual Action Flea & Tick Collar year round for most effective control.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Do not open until ready to use. Dust will form on Collar during storage. Do not get dust in mouth or eyes. Harmful if swallowed. If in eyes will cause eye irritation. In case of contact, flush eyes with water. If irritation persists, get medical attention. Collar is intended for external use only, not to be taken internally. Some animals may become irritated by any Collar. If this occurs, remove Collar. If conditions persists, consult a veterinarian.


NOTE TO PHYSICIAN/VETERINARIAN: Dust released by this Collar is a cholinesterase inhibitor. Atropine is antidotal.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Place Collar around pet's neck, adjust for fit, and buckle in place. The Collar must be worn loosely to allow for growth of the pet and to permit the Collar to move about the neck. Generally, a properly fitted Collar is one that when fastened, will snugly slide over pet's head. Leave 2 or 3 inches for extra adjustment. Cut off and dispose of excess length. Wash hands after installing Collar on pet. Under normal conditions, replace Collar every 6 months. Under conditions of severe infestation and where continued rapid kill is desired, collar may be replaced more often. Do not use on pet's under 6 weeks of age.

STORAGE AND DISPOSAL: Store unopened Collar in a cool dry place. Dispose of empty pouch and expired Collar in trash

Distributed by:
ConAgra Pet Products
Omaha, NE 68102



EPA Reg. No. 2517-61
EPA Est. No. 2517-VA-1

Made in USA

UPC

Contact Info/Guarranty/Offer