

2517-129

09/05/2013

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 05, 2013

Mr. James Messina
Authorized Representative of Sergeant's Pet Care Product, Inc.
c/o Exponent
1150 Connecticut Ave. NW, Suite 1100
Washington, DC 20036

Subject: Application for Pesticide Amendment
Cyphenothrin + Pyriproxyfen Squeeze-On for Dogs (20%)
EPA Reg. No. 2517-129
Your Submission Dated February 26, 2013
Decision No: 480880

Dear Mr. Messina:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

Conditions of acceptance

1. This registration is now time-limited and expires two years from the date of this letter.
2. Please consider limiting the formulations of this product to one basic confidential statement of formula. The Agency may require this in the future and no additional alternate formulations or minor formulation amendments will be approved for this product in the future.
3. You must submit quarterly enhanced incident reports and quarterly sales information in doses sold for this product beginning January 1, 2014. Please flag any Confidential Business Information as such. Enhanced incident reporting should be submitted to the Product Manager. Please flag any Confidential Business Information as such. Enhanced incident reporting should be submitted to the Product Manager. Quarterly sales information should be submitted to the attention of Kaitlin Keller, Pesticide Reevaluation Division, Mail Code 7508P.

The following is a list of information that must be included in the quarterly reports for each incident:

EPA Registration Number
Product name (brand name)
Lot #
Where purchased: internet, store, veterinarian
Active Ingredient(s)
Weight range for product

Date on which incident occurred. (mm/dd/yyyy)
State in which the incident occurred. (standard 2 letter abbreviation)
Registrant case #

Species: dog, cat, other (specify)
Breed: (as reported by pet owner)
Age: months or years
Sex: M, F, or neutered
Weight: pounds

Primary Route of Exposure: dermal, oral, other animal, inhalation, other
Body System: neurological, dermatological, GI, respiratory, ocular, other
Major signs noted with separate column for each sign, using standard terminology
Time to Onset: (hours, days)
Treated by veterinarian: yes or no
First time product used: yes or no
Misuse: use on incorrect species, overdose, too frequent dosing, other (describe)
Any known precondition
EPA Severity Code: death, major, moderate, minor
Outcome: died, recovered, still treated, unknown


4. Along with the enhanced incident reporting, you must submit an analysis of the incidents seen, to include the following details:

- All incidents should be reported including all minor dermal and ocular irritation reports.
- Summary table for dogs showing number of incidents of each severity code for each route of exposure. Each incident should only be reported once. If one incident has several routes of exposure, the order should be ocular > oral > dermal. In other words, an incident with both oral and dermal exposure would be reported as oral exposure, and an incident with both ocular and oral exposure would be reported as ocular exposure.

- A similar summary table for cats (misuse or secondary exposure) showing number of incidents of each severity code for each route of exposure.
- Summary table for dogs and table for cats showing number of incidents that are believed due to secondary exposure (e.g., multi-pet households).
- A summary table for dogs showing number of incidents for each severity code for these age ranges: <3 months, 3-6 months, 6-9 months, 9-12 months, 1 yr, 2 yr, 3 yr, 4 yr, 5 yr, 6 yr, 7 yr, 8 yr, 9 yr, 10 yr, 11 yr, 12 yr, 13 yr, 14 yr, 15 yr, >15 yr.
- A summary table showing the number of dog incidents for each severity code for each pet weight range on the product label.
- A summary table for dog weight showing number of incidents for each product weight range. This table should show number of incidents in dogs weighing less than that product weight range, number of incidents in dogs in lower half of weight range, number of incidents in dogs in upper half of weight range, and dogs weighing more than the product weight range.
- Table showing number of incidents for each dog breed.
- Table showing number of incidents in dogs for each clinical sign.
- Table showing number of incidents in dogs for each organ system.
- Report aggregate incidents, but do not combine moderate and minor incidents.

If EPA determines that future mitigation measures are necessary for all spot-on products, the Agency will inform registrants and if needed, EPA may take regulatory action. If you have any questions contact Driss Benmhend at 703-308-9525 or by e-mail at: Benmhend.driss@epa.gov.

Sincerely,



Mark Suárez
Product Manager 13, Insecticide Branch
Registration Division (7504P)

Enclosure: *Label stamped "Accepted"*

Sergeant's Master Label

4/11

03 Sep 2013 Mitigation
Information in [] is optional.

v13.1
Page 1 of 8

EPA No. 2517-129
EPA SAL 09 Oct 2012

FRONT PANEL DOG SQUEEZE ON Cyphenothrin & Pyriproxyfen

[Optional text appears in brackets – the final label may include some or all of the optional text on front, back or side label panels]

Cyphenothrin+Pyriproxyfen Squeeze-On for Dogs (20%)

[ABNs:]

[Sergeant's [Plus] [Flea and Tick] Squeeze-On for Dogs [and Puppies Over 12 Weeks Old]]

[Sergeant's Gold® [Plus] [Flea and Tick] Squeeze-On for Dogs [and Puppies Over 12 Weeks Old]]

[SentryPro XFC [Flea and Tick] Squeeze-On for Dogs [and Puppies Over 12 Weeks Old]]

[Sergeant's Evolve [11] [21] [40] Flea and Tick Squeeze-On for Dogs]

[Sentry®Pro XTR [11] [21] [40] Flea and Tick Squeeze-On for Dogs]

[Sentry®Pro XFT [11] [21] [40] Flea and Tick Squeeze-On for Dogs]

[Sentry®Pro XFTC [11] [21] [40] Flea and Tick Squeeze-On for Dogs]

[Sentry®Pro XFC+ [11] [21] [40] Flea and Tick Squeeze-On for Dogs]

[1.5cm x 1.5 cm on final or as indicated]



[For use ONLY on dogs 9 - 20 lbs and 12 weeks of age or older]

[For use ONLY on dogs [21 - 39 lbs] [40 - 60 lbs]] {OR}

[Only for use on dogs over 12 weeks of age and 9 lbs to 20 lbs]

[Only for use on dogs [21 lbs to 39 lbs] [40 lbs to 60 lbs]]

ACCEPTED

SEP - 5 2013

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

[Specially Formulated for Dogs and Puppies (over 12 weeks old)]

[Flea & Tick Control for Dogs & Puppies 12 weeks old and older]

[For Dogs & Puppies Over 12 Weeks of Age [greater than 9 lbs] [9 to 20 lbs] [21 to 39 lbs] [40 to 60 lbs]]

EPA Reg. No: 2517-129

[[12][15][18][21][24][32][52] Week Supply Only For Use on Dogs weighing [more than 9 lbs], [9 to 20 lbs], [21 to 39 lbs], [40 to 60 lbs]]

[Apply once every [4 weeks] [1 month] [30 days]!]

[Up to [4 week] [1 Month] Dose!]

[(ICON Showing up to 4 Week Protection)]

[Three Applications {for cartons with 3 applicators}] and/or [3 Month Supply] or [12 Week Supply]

[Four Applications {for cartons with 4 applicators}] and/or [4 Month Supply] or [16 Week Supply]

[Best if used year round!]

[Easy to Use Application]

[[Free] [Brochure] [and] [Calendar] [Stickers] Inside!]

[Patent pending technology] [U.S. and Foreign patent pending technology]

[Dogs can be bathed 24 hours after squeeze-on is applied.]

[Quick Drying] [formula]

[with Vitamin E] [No oily residue]

[Fragrance Free]

[Formulated] [with] DERMAshield™ [Technology]

[DERMAshield™ [Technology] [which] dries quickly to keep the product on the outer skin surfaces]

[and contains [a vitamin] [vitamin E] to soothe skin]]

[Not all dogs are the same. Should their flea and tick [control][product] be?]

[Brand Name][developed for the weight of your dog][specially formulated for the weight of your dog]

Sergeant's Master Label

03 Sep 2013 Mitigation
Information in [] is optional.

v13.1
Page 2 of 8

EPA No. 2517-129
EPA SAL 09 Oct 2012

51/11

[Multiple species]

[Kills fleas, ticks, flea eggs, flea larvae][Repels mosquitoes]
[Kills fleas, ticks][Kills flea eggs, flea larvae] [Repels mosquitoes]
[[Four] [4] Way Protection! [Kills ticks, fleas, flea eggs & flea larvae]]
[[Five] [5] Way Protection! [Kills fleas, flea eggs, flea larvae, and ticks, and repels mosquitoes]]
[Five in one protection [Kills fleas, flea eggs, flea larvae, and ticks, and repels mosquitoes]]
[Kills all stages of the flea life cycle]
[Kills fleas, ticks, flea eggs, and flea larvae [and repels mosquitoes,]]
[Kills fleas, flea eggs, flea larvae, ticks, and repels mosquitoes]
[Kills fleas [in as little as 1 hour], [and] [ticks] [in as little as 3 hours], [flea eggs & flea larvae] [and repels mosquitoes]]
[Kills Fleas [in as little as 1 hour] [and] Ticks [in as little as 3 hours]]
[Kills Fleas, Ticks, Flea Eggs & Flea Larvae [and repels Mosquitoes] for up to [4 Weeks] [1 month]]
[Works for up to 4 Weeks on fleas and ticks!]
[Protects for up to 4 Weeks on [adult] fleas and ticks!]
[Up to 4 Week [One Month] [Adult] Flea and Tick [Control] [Protection!] [Treatment!]]
[Kills [Adult] Fleas and Ticks for up to 4 weeks!]
[Starts killing fleas and ticks in [less than][just] [5][15] minutes] [after application]
[Starts to kill fleas and ticks in [just] [5][15] minutes] [after application]
[Stops][Prevents] fleas [from biting][before they bite]
[Kills fleas before they bite]
[Kills fleas in [less than] [just] 1 hour [after application]]
[Quickly kills fleas in [just] [1 hour] and ticks [in as little as 3 hours]]
[Kills fleas [as quick as] [just] 1 hour [after application] and ticks in [as quick as] [just] 3 hours [after application]]
[Kills Fleas, Ticks] [Kills Flea Eggs, Flea larvae][Repels mosquitoes]

[Flea and Flea eggs and Larval]

[Starts killing fleas in [less than][just] [5][15] minutes] [after application]
[Starts to kill fleas in [less than] [just] [5][15] minutes] [after application]
[Kills fleas in 1 hour] [after application]
[Stops][Prevents] fleas [from biting][before they bite][dogs][your dog]
[Stops][Prevents] fleas from ingesting a blood meal
[Stops fleas from biting [which may cause anemia]
[Stops fleas from biting [which may cause [flea allergy] dermatitis]
[[Stops][Kills] fleas which may cause flea allergy dermatitis]
[[Stops][Kills] fleas which may cause anemia]
[Kills fleas before they [can] bite]
[Quickly kills fleas in [less than] [just] [1 hour]]
[With Pyriproxyfen to Break Flea Life Cycle]
[With Insect Growth Regulator to Break Flea Life Cycle]
[With [Pyriproxyfen] Insect Growth Regulator which Breaks the Flea Life Cycle]
[Dual Action [!]: Effectively Breaks the Flea Life Cycle [!]]
[Dual Action [!]: Cyphenothrin with [Pyriproxyfen] IGR Effectively Breaks the Flea Life Cycle]
[Prevents Flea Eggs From Developing Into Biting Adults]
[[Breaks Flea Life Cycle!] for up to 9 weeks]
[Controls reinfestation for up to 9 weeks!]
[Kills Flea Eggs & Flea Larvae for up to [63 days], [9 weeks]!]
[Kills for up to [9 weeks] [63 days] [flea eggs][flea larvae]]

Sergeant's Master Label

6/11

03 Sep 2013 Mitigation
Information in [] is optional.

v13.1
Page 3 of 8

EPA No. 2517-129
EPA SAL 09 Oct 2012

[Kills fleas in as little as 1 hour!]
[Kills Fleas for up to 4 weeks!]
[Kills Adult Fleas for up to 4 weeks!]

[Ticks]

[Starts killing ticks in [less than][just] [5][15] minutes] [after application]
[Starts to kill ticks in [just] [5][15] minutes] [after application]
[Kills ticks in 3 hours] [after application]
[Quickly kills ticks in [as little as] [just] 3 hours][!]
[Kills and Detaches Ticks]
[Kills ticks in [as little as][just] 3 hours][!]
[Prevents ticks from attaching and feeding[in as little as][3 hours]]
[Kills ticks including those that may transmit Lyme disease]
[Kills Ticks for Up to [4 weeks] [1 month] [30 days]!]
[Kills Deer Ticks (vector of Lyme disease) for up to 4 weeks]
[Kills deer ticks [before they can] [that] [transmit Lyme disease,] [for up to [4 weeks] [1 month] [30 days]]]
[Kills Ticks (Including Deer Ticks) for up to [4 weeks] [1 month] [30 days]]
[Kills [brown dog ticks] [(*Rhipicephalus sanguineus*)] [for up to [4 weeks] [1 month] [30 days]]]
[Kills American Dog Ticks [(*Dermacentor variabilis*)] for up to [4 weeks] [1 month] [30 days]!]

[Mosquitoes]

[Repels mosquitoes]
[Repels mosquitoes within 24 hours]
[Repels mosquitoes] [for up to [30 days] [4 weeks] [1 month] !]
[Repels mosquitoes that may transmit heartworm]
[Repels mosquitoes that may transmit [heartworm] [West Nile virus] [from feeding on dogs] [for up to [one month] [30 days][4 weeks]]
[Repels Mosquitoes] [Prevents mosquitoes that transmit West Nile virus and heartworm] [from feeding on dogs] [for up to [one month] [30 days][4 weeks]]
[Prevents mosquitoes from blood feeding, which may transmit viruses such as West Nile Virus and heartworm to dogs for up to four weeks.]

ACTIVE INGREDIENTS:

Cyphenothrin (CAS# 39515-40-7)20.00%
Pyriproxyfen (CAS# 95737-68-1) 2.00%

OTHER INGREDIENTS[*]: 78.00%

[TOTAL] 100.00%

[*(Diethylene glycol monoethyl ether, vitamin E nicotinate)]

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See [Back][or][Side][Label[s]][Panel[s] for Additional Precautionary Statements

Sergeant's Master Label

03 Sep 2013 Mitigation
Information in [] is optional.

v13.1
Page 4 of 8

EPA No. 2517-129
EPA SAL 09 Oct 2012

7/11

NET CONTENTS: XXX fl oz (XXX mL)

[THREE][FOUR][SIX][TWELVE]

[0.05 fl oz (1.5 mL) Tubes] [0.10 fl oz (3.0 mL) Tubes] [0.15 fl oz (4.5 mL) Tubes]

{END – FRONT PANEL}

BACK/SIDE PANEL DOG SQUEEZE ON Cyphenothrin & Pyriproxyfen

[Apply Once Every Month] [Can be applied monthly [if needed]]

**READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS AND PUPPIES OVER 12 WEEKS OF AGE.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if swallowed or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. If your dog is exhibiting signs of and/or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

FIRST AID

IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF SWALLOW ED	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Have person sip a glass of water if able to do so.• Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-781-4738.

NOTE TO PHYSICIAN OR VETERINARIAN

Treat patient symptomatically

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR

Sergeant's Master Label

03 Sep 2013 Mitigation
Information in [] is optional.

v13.1
Page 5 of 8

EPA No. 2517-129
EPA SAL 09 Oct 2012

DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on cats or animals other than dogs. Do not allow your dog to ingest this product. Do not use on puppies under 12 weeks of age or less than 9 lbs. Use entire contents of tube vial on each dog. [Do not split one tube between dogs.] [Do not use multiple tubes on one dog.] Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

Side Effects: Monitor your dog after application. [Some dogs may experience temporary startle effects when any product is applied.] Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-800-781-4738. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.

[1.5cm x 1.5cm on final or as indicated]



DO NOT USE ON CATS. MAY BE FATAL. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian immediately.

{Outer Packaging}

How to open: Remove product tube[s] [vial] from the package. [Separate one tube from the others.] [Holding][Hold] the tube [vial] with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.] **OR** [Holding][Hold] the tube [vial] with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.]

How to apply: Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Repeat application may be made if necessary, but do not apply more often than once every 4 weeks.

[Optional drawings that may or may not appear anywhere on finished label]



Frequency of Application:

[When used monthly, {product name} breaks the flea life cycle and controls flea and tick infestations.]

Only For Dogs Weighing Between 9 and 20 lbs

{For cartons containing one 1.5 mL (0.05 fl oz) applicator tubes}

Apply one tube (1.5 mL)(0.05 fl oz) [as a spot to the dog's back between the shoulder blades]or [from the back of the neck to a point midway between the neck and tail]. Do not apply to dogs weighing less than 9 lbs.

Sergeant's Master Label

03 Sep 2013 Mitigation
Information in [] is optional.

v13.1
Page 6 of 8

EPA No. 2517-129
EPA SAL 09 Oct 2012

9/11

Only For Dogs Weighing Between 21 and 39 lbs

{For cartons containing two 1.5 mL (0.05 fl oz) applicator tubes}

[Apply two tubes (1.5 mL)(0.05 fl oz) as a spot on the dog's back between the shoulder blades] [from the back of the neck to a point midway between the neck and tail].

OR

{For cartons containing one 3.0 mL (0.10 fl oz) applicator tube}

[Apply one tube (3.0 mL)(0.10 fl oz) as a spot or stripe to the dog's back between the shoulder blades.]
or [from the back of the neck to a point midway between the neck and tail.]

Only For Dogs Weighing Between 40 and 60 lbs

{For cartons containing three 1.5 mL (0.05 fl oz) applicator tubes}

[Apply two tubes (1.5 mL)(0.05 fl oz) as a spot or stripe to the dog's back between the shoulder blades and apply the third tube from the back of the neck to a point midway between the neck and tail].

OR

{For cartons containing one 4.5 mL (0.15 fl oz) applicator tube}

[Apply one tube (4.5 mL)(0.15 fl oz) from the back of the neck to a point midway between the neck and tail]. Do not use two applicator tubes on dogs larger than 60 lbs.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F [25°C]), dry place out of reach of children. Do not refrigerate. Protect from direct sunlight.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

[Brand name] is an effective and easy to use product.] [Brand name] prevents eggs from fleas on treated dogs from developing into biting adults for up to 63 days] [9 weeks]. [As with all flea and tick control products, [Brand name] should be used as part of a program aimed at reducing flea populations in the dog's environment (bedding, carpets, kennel, yard).] [Consult your retailer for program recommendations.]

[WARRANTY

Read the entire Directions for Use and Warranty before using this Product. By using this product, user or buyer accepts the following warranty. The directions for use of this product are believed to adequate and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Unintended consequences may result because of unknown factors. All such risks shall be assumed by the user or buyer. Sergeant's is committed to providing high quality products. To the extent consistent with applicable law, Sergeant's makes no warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. To the extent consistent with applicable law, Sergeant's, the manufacturer, or the Seller shall not be liable for indirect, special, incidental or consequential damages. To the extent consistent with applicable law, the exclusive remedy of the user or buyer shall not exceed the purchase price paid.]

[If you have questions or comments about this product, please write: [Sergeant's] [SENTRY]
Consumer Response, P.O. Box 540399, Omaha, NE 68154-0399]

Sergeant's Master Label

10/11

03 Sep 2013 Mitigation
Information in [] is optional.

v13.1
Page 7 of 8

EPA No. 2517-129
EPA SAL 09 Oct 2012

In Case of Emergency, call: 1-800-781-4738 [Coupon(s) inside or on outside of box]
Non-Emergency, call: 1-800-224-PETS (7387) [Made in USA][Manufactured in USA]

Made in the USA For:] or [Manufactured by:][Distributed by:]


Sergeant's Pet Care Products, Inc.
Omaha, NE 68138-3710

EPA Reg. No. 2517-129
EPA Est. No. XXXXX-XX-XX
[lot, date & label code] [UPC CODE]

{END - BACK / SIDE PANEL}

OUTER PACKAGING OR FOLDED PANEL/FLAP OF FRONT OF PACKAGING DOG SQUEEZE ON Cyphenothrin & Pyriproxyfen

[Optional drawings and information that may or may not appear anywhere on finished label]



(Insert diagram that includes checklist with magnifying glass and checklist graphic)

[Look at the Label (Look at the Label icon) [Easy] 6 Step [Easy] [Spot On] [Squeeze-On] [Topical] Checklist]

- [1] Read the label completely and follow directions
- [2] Weigh your dog
- [3] Do NOT use dog products on cats
- [4] Do NOT treat with more than one pesticide product at a time
- [5] Separate animals after [treatment][application] for at least 24 hours to avoid chance of ingestion
- [6] Do NOT split tubes between dogs [and use the ENTIRE tube contents.]

[Brand Name]

[1][2][3][4][5][6][12][24][Month [Application] Tracker]

[For year-round protection, apply [Brand Name] monthly.]

[Dog's Name (Empty blank for owner to fill in name)]

[First][1st][Second][2nd][Third][3rd][Fourth][4th][Fifth][5th][Sixth][6th][Seventh][7th][Eighth][8th][Ninth][9th][Tenth][10th][Eleventh][11th][Twelfth][12th]

[Dose] [Application](Empty blank for owner to fill in date)(Date: MM/DD/YY)

[calendar stickers with [Brand Name]

[Place stickers on your calendar as a reminder to reapply [Brand Name] [in 30 days.]

[How to open: Remove product tube[s] [vial] from the package. [Separate one tube from the others.] [Holding][Hold] the tube [vial] with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.] OR [Holding][Hold the tube [vial] with notched end

Sergeant's Master Label

03 Sep 2013 Mitigation
Information in [] is optional.

v13.1
Page 8 of 8

EPA No. 2517-129
EPA SAL 09 Oct 2012

pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.]]

[How to apply: Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Repeat application may be made if necessary, but do not apply more often than once every 4 weeks.]

[Only For Dogs Weighing Between 9 and 20 lbs

{For cartons containing one 1.5 mL (0.05 fl oz) applicator tubes}

Apply one tube (1.5 mL)(0.05 fl oz) [as a spot to the dog's back between the shoulder blades]or [from the back of the neck to a point midway between the neck and tail].] Do not apply to dogs weighing less than 9 lbs.

[Only For Dogs Weighing Between 21 and 39 lbs

{For cartons containing two 1.5 mL (0.05 fl oz) applicator tubes}

[Apply two tubes (1.5 mL)(0.05 fl oz) as a spot on the dog's back between the shoulder blades] [from the back of the neck to a point midway between the neck and tail].

OR

{For cartons containing one 3.0 mL (0.10 fl oz) applicator tube}

[Apply one tube (3.0 mL)(0.10 fl oz) as a spot or stripe to the dog's back between the shoulder blades.] or [from the back of the neck to a point midway between the neck and tail.]]

[Only For Dogs Weighing Between 40 and 60 lbs

{For cartons containing three 1.5 mL (0.05 fl oz) applicator tubes}


[Apply two tubes (1.5 mL)(0.05 fl oz) as a spot or stripe to the dog's back between the shoulder blades and apply the third tube from the back of the neck to a point midway between the neck and tail].

OR

{For cartons containing one 4.5 mL (0.15 fl oz) applicator tube}

[Apply one tube (4.5 mL)(0.15 fl oz) from the back of the neck to a point midway between the neck and tail].] Do not use two applicator tubes on dogs larger than 60 lbs.

[Optional graphics that may or may not appear anywhere on finished label]




{END - BACK / SIDE PANEL}

TUBE / VIAL LABEL DOG SQUEEZE ON Cyphenothrin & Pyriproxyfen

Front Label

Back Label

 <p>{Brand Name} For Dogs [9 to 20][21-39][40-60] lbs ≥12 wks Cyphenothrin 20.00%, Pyriproxyfen 2.00% 0.05 fl oz (or) 0.10 fl oz (or) 0.15 fl oz {label code}</p>	<p>KEEP OUT OF REACH OF CHILDREN CAUTION: Read directions and precautions before using. Use scissors to open. EPA REG. No. 2517-129 {label code}</p>
---	--

{END - TUBE/VIAL COPY}