



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 20, 2017

Prince Bamaze
Regulatory Affairs
Sergeant's Pet Care Products Inc.
10077 South 134th Street
Omaha, NE 68138-3710

Subject: Label Amendment – Agency Initiated Master Label amendment to address safety mitigation
Product Name: Cyphenothrin + Pyriproxyfen Squeeze-On for Dogs (20%)
EPA Registration Number: 2517-129
Application Date: 10/02/2017
Decision Number: 534894

Dear Mr. Bamaze:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 2517-129
Decision No. 534894

with FIFRA section 6. If you have any questions, please contact Helen Hull-Sanders by phone at 703-347-0243, or via email at hull-sanders.helen@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Elizabeth Fertich". The signature is written in a cursive style with a large, looped initial "E".

Elizabeth Fertich
Product Manager 4
Invertebrate & Vertebrate Branch 1
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 1 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

FRONT PANEL

DOG SQUEEZE ON Cyphenothrin & Pyriproxyfen

[Optional text appears in brackets – the final product label may include some or all of the optional text on front, back or side label panels. The CRP required text and format will not change from current label.]

{Information is instructional}

Cyphenothrin+Pyriproxyfen Squeeze-On for Dogs (20%)

{Market Label - the word 'Dog' will be 40-75% in height of the largest letter in the primary brand name.}

{Market Label-a large clear picture of a Dog in the appropriate weight range will be on the front panel of the label}

{A minimum of 1.5 cm x 1.5 cm icon as indicated black cat, black text, red circle and line through cat, on white background to ensure easy spotting, in lower right-hand corner of front label panel.}



{One of the statements below will appear prominently on front panel of market label as required}

For use ONLY on [Small][Medium][Large] dogs [&] [and] puppies [weighing] [9-20] [21-39] [40-60] lbs. [over] 12 weeks [of age] or older.

{OR}

ONLY for [Small][Medium][Large] dogs [&] [and] puppies [weighing] [9-20] [21-39] [40-60] lbs. [over] 12 weeks [of age] or older.

ACTIVE INGREDIENTS:

Cyphenothrin [(CAS# 39515-40-7)]20.00%

Pyriproxyfen (CAS# 95737-68-1)2.00%

OTHER INGREDIENTS [*]:78.00%

TOTAL100.00%


[*(Diethylene glycol monoethyl ether, vitamin E nicotinate)]

KEEP OUT OF REACH OF CHILDREN CAUTION

See [Back][Side][Label(s)][Panel(s)] [or][Insert] for First Aid [and] [Additional] Precautionary Statements

NET CONTENTS:

- (1) [0.05 fl oz (1.5 mL) Tube(s) [Vial(s)]]
[0.10 fl oz (3.0 mL) Tube(s) [Vial(s)]]
[0.15 fl oz (4.5 mL) Tube(s) [Vial(s)]]
{OR}
- (2) [ONE] [0.05 fl oz (1.5 mL) Tube [Vial]]
[TWO] [0.05 fl oz (1.5 mL) Tubes [Vials]]
[THREE] [0.05 fl oz (1.5 mL) Tubes [Vials]]



{END – FRONT PANEL}

Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 2 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

BACK/SIDE PANEL

DOG SQUEEZE ON Cyphenothrin & Pyriproxyfen

[Apply Once Every Month]

[Can be applied monthly]

**READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS AND PUPPIES OVER 12 WEEKS OF AGE.
[WEIGHING] [9-20], [21-39], [40-60] lbs.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if swallowed or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS: For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. This product is not a substitute for prescribed heartworm preventative. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged, pregnant, nursing dogs, or dog with known organ dysfunction. If your dog is exhibiting signs of and/or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Have person sip a glass of water if able to do so.• Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-781-4738.	
NOTE TO PHYSICIAN OR VETERINARIAN	
Treat patient symptomatically	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT ALLOW CHILDREN TO APPLY PRODUCT. To prevent harm to you and your dog, read entire label before each use. Follow all directions and precautionary statements carefully. Dog must be dry at time of application and remain dry for 24 hours after application. Use entire contents of [tube] [vial] on each dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 3 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

RESTRICTIONS:

Do not use on cats or animals other than dogs.

Do not allow your dog to ingest this product.

Do not use on puppies under 12 weeks of age or less than 9 lbs. [21 lbs.] [40 lbs.].

Do not bathe or allow dog to swim for 24 hours after application.

Do not apply more than one [tube][vial] per treatment for one dog regardless of dog's weight. **{OR}**

Do not apply more than contents of one box per treatment to one dog regardless of dog's weight.

Do not reapply for [30 days][4 weeks][one month].

{Side Effects box to be located immediately above the cat warning box on the back label panel}

Side Effects: Monitor your dog after application. [Some dogs may experience temporary startle effects when any product is applied.] Dogs may experience [some] temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If any side effects occur or your dog has unusual reaction to the application, consult your veterinarian or call 1-800-781-4783. Consult your veterinarian before repeating application.

{A minimum of 1.5cm x 1.5cm cat prohibition icon and warning text will be in a box in the lower right hand corner of the back panel with black cat + red circle and line through the cat, black text, on white background to ensure easy spotting.}

DO NOT USE ON CATS. MAY BE FATAL. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian immediately.




[See Insert inside [or] [Folded Panel(s)/Flap] for information on **How to open** and **How to apply**]

{Referral Statement used only when 'Directions for Use' on outer packaging does not contain information below}

How to open: Remove product tube(s) [vial(s)] from the package. [Separate one tube [vial] from the others.] [Holding][Hold] the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.] **OR** [Holding] [Hold] the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.]

How to apply: Invert tube [vial] over dog and use open end to part dog's hair. Squeeze tube [vial] firmly to apply all of the solution to the dog's skin, as directed below. Repeat application if necessary, but not more than once every 4 weeks.

[Optional drawing that may or may not appear anywhere on finished label]



Frequency of Application:

[When used monthly, {product name} breaks the flea life cycle and controls flea and tick infestations.]

Only For Dogs Weighing Between 9 and 20 lbs

{Cartons containing one 1.5 mL (0.05 fl oz) tube [vial]}

Apply one tube[vial] (1.5 mL)(0.05 fl oz) [as a spot to the dog's back between the shoulder blades] {or} [from the back of the neck to a point midway between the neck and tail].

Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 4 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

Only For Dogs Weighing Between 21 and 39 lbs

{(1) - Cartons containing one 3.0 mL (0.10 fl oz) tube[vial]}

Apply one tube [Vial] (3.0 mL)(0.10 fl oz) [as a spot or stripe to the dog's back between the shoulder blades] {or} [from the back of the neck to a point midway between the neck and tail.]

{OR}

{(2) - Cartons containing two 1.5 mL (0.05 fl oz) tubes[vials]}

Apply two tubes [Vial] (1.5 mL)(0.05 fl oz) [as a spot on the dog's back between the shoulder blades] {or} [from the back of the neck to a point midway between the neck and tail].

Only For Dogs Weighing Between 40 and 60 lbs

{(1) - Cartons containing one 4.5 mL (0.15 fl oz) tube [Vial]}

Apply one tube [Vial] (4.5 mL)(0.15 fl oz) from the back of the neck to a point midway between the neck and tail.

{OR}

{(2) - Cartons containing three 1.5 mL (0.05 fl oz) tubes[vials]}

Apply two tubes [vial] (1.5 mL)(0.05 fl oz) as a spot or stripe to the dog's back between the shoulder blades and apply the third tube from the back of the neck to a point midway between the neck and tail.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube [vial] from the pack until ready to use. Store in a cool (below 77°F [25°C]) dry place out of reach of children. Do not refrigerate. Protect from direct sunlight.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain

[[Brand name] is an effective and easy to use product. [Brand name] prevents eggs from fleas on treated dogs from developing into biting adults for up to [63 days] [9 weeks]. [As with all flea and tick control products, [Brand name] should be used as part of a program aimed at reducing flea populations in the dog's environment (bedding, carpets, kennel, yard).] [Consult your retailer for program recommendations.]]

[WARRANTY

Read the entire Directions for Use and Warranty before using this Product. By using this product, user or buyer accepts the following warranty. The directions for use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Unintended consequences may result because of unknown factors. All such risks shall be assumed by the user or buyer. Sergeant's is committed to providing high quality products. To the extent consistent with applicable law, Sergeant's makes no warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. To the extent consistent with applicable law, Sergeant's, the manufacturer, or the Seller shall not be liable for indirect, special, incidental or consequential damages. To the extent consistent with applicable law, the exclusive remedy of the user or buyer shall not exceed the purchase price paid.]

[If you have questions or comments about this product, please write: [Sergeant's] [SENTRY] Consumer Response, P.O. Box 540399, Omaha, NE 68154-0399.]

In Case of Emergency, call: 1-800-781-4738 [Coupon(s) inside or on outside of box]
Non-Emergency call 1-800-224-PETS (7387) {When outside of the box, coupon will not cover or distract consumer from required text}

Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 5 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

[Made in USA][Manufactured in USA]
[Made in the USA For:] or [Manufactured by:][Distributed by:]


Sergeant's Pet Care Products, Inc.
Omaha, NE 68138-3710

EPA Reg. No. 2517-129
EPA Est. No. XXXXX-XX-XX
[lot, date & label code] [UPC CODE]

{END - BACK / SIDE PANEL}

TUBE / VIAL LABEL DOG SQUEEZE ON Cyphenothrin & Pyriproxyfen

Front Label

{Brand Name} 
For Dogs [9 to 20][21-39][40-60] lbs \geq 12 wks
Cyphenothrin 20.00% , Pyriproxyfen 2.00%
0.05 fl oz (**or**) 0.10 fl oz (**or**) 0.15 fl oz

Back Label

KEEP OUT OF REACH OF CHILDREN
CAUTION: Read entire label before each use.
Use scissors to open.
EPA Reg. No. 2517-129
{Lot No.} {label code}


{Lot No. is engraved on the crimp portion of the tube} {label code refers to an internal product tracking code}
{EPA Reg. No. may be on either the front or back label}

{END - TUBE / VIAL COPY}

INSERT OR FOLDED PANEL/FLAP OF FRONT OF PACKAGING DOG SQUEEZE ON Cyphenothrin & Pyriproxyfen

**READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS AND PUPPIES OVER 12 WEEKS OF AGE.
[WEIGHING] [9-20], [21-39], [40-60] lbs.**

[Optional graphics below may or may not appear anywhere on finished label]



Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 6 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

(Insert diagram that includes checklist with magnifying glass and checklist graphic)

[Look at the Label (Look at the Label icon) [Easy] 6 Step [Easy] [Spot On] [Squeeze-On] [Topical] Checklist]

- [1] Read the label completely and follow directions
- [2] Weigh your dog
- [3] Do NOT use dog products on cats
- [4] Do NOT treat with more than one pesticide product at a time
- [5] Separate treated dog from other pets for at least 24 hours to avoid chance of ingestion
- [6] Do NOT split tubes [vials] between dogs [and use the ENTIRE tube [vial] contents.]

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Have person sip a glass of water if able to do so.• Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-781-4738.	
NOTE TO PHYSICIAN OR VETERINARIAN	
Treat patient symptomatically	

[Brand Name]

[1][2][3][4][5][6][12][24][Month [Application] Tracker]

[For year-round protection, apply [Brand Name] monthly.]

[Dog's Name (Empty blank for owner to fill in name)]

[First][1st][Second][2nd][Third][3rd][Fourth][4th][Fifth][5th][Sixth][6th][Seventh][7th][Eighth][8th][Ninth][9th][Tenth][10th][Eleventh][11th][Twelfth][12th]

[Dose] [Application](Empty blank for owner to fill in date)(Date: MM/DD/YY)

[Calendar stickers with [Brand Name]

[Place stickers on your calendar as a reminder to reapply [Brand Name] [in 30 days.]

How to open: Remove product tube(s)[vial(s)] from the package. [Separate one tube [vial] from the others.] [Holding][Hold] the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.] **OR** [Holding] [Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.]

Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 7 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

How to apply: Invert tube [vial] over dog and use open end to part dog's hair. Squeeze tube [vial] firmly to apply all of the solution to the dog's skin, as directed below. Repeat application if necessary, but not more than once every 4 weeks.



Frequency of Application:

[When used monthly, {product name} breaks the flea life cycle and controls flea and tick infestations.]

Only For Dogs Weighing Between 9 and 20 lbs

{For cartons containing one 1.5 mL (0.05 fl oz) tube [vial]}

Apply one tube [vial] (1.5 mL)(0.05 fl oz) [as a spot to the dog's back between the shoulder blades.] {or} [from the back of the neck to a point midway between the neck and tail]. **Do not apply more than one [tube][vial] per treatment for one dog regardless of dog's weight.**

Only For Dogs Weighing Between 21 and 39 lbs

{(1) - Cartons containing one 3.0 mL (0.10 fl oz) tube[vial]}

[Apply one tube [Vial] (3.0 mL)(0.10 fl oz) [as a spot or stripe to the dog's back between the shoulder blades.] {or} [from the back of the neck to a point midway between the neck and tail.] **Do not apply more than one [tube][vial] per treatment for one dog regardless of dog's weight.**

{OR}

{(2) - Cartons containing two 1.5 mL (0.05 fl oz) tubes[vials]}

[Apply two tubes [Vial] (1.5 mL)(0.05 fl oz) [as a spot on the dog's back between the shoulder blades] {or} [from the back of the neck to a point midway between the neck and tail]. **Do not apply more than contents of one box per treatment to one dog regardless of dog's weight.**

Only For Dogs Weighing Between 40 and 60 lbs

{(1) - Cartons containing one 4.5 mL (0.15 fl oz) tube [Vial]}

Apply one tube [Vial] (4.5 mL)(0.15 fl oz) from the back of the neck to a point midway between the neck and tail. **Do not apply more than one [tube][vial] per treatment for one dog regardless of dog's weight.**

{OR}

{(2) - Cartons containing three 1.5 mL (0.05 fl oz) tubes[vials]}

Apply two tubes [vial] (1.5 mL)(0.05 fl oz) as a spot or stripe to the dog's back between the shoulder blades and apply the third tube from the back of the neck to a point midway between the neck and tail.

Do not apply more than contents of one box per treatment to one dog regardless of dog's weight.


Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 8 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

[Optional graphics below may or may not appear anywhere on finished label]



{END - INSERT OR FOLDED PANEL/FLAP}

[ABNs:]

[Sergeant's [Plus] [Flea and Tick] Squeeze-On for Dogs [and Puppies Over 12 Weeks Old]]
[Sergeant's Gold® [Plus] [Flea and Tick] Squeeze-On for Dogs [and Puppies Over 12 Weeks Old]]
[SentryPro XFC [Flea and Tick] Squeeze-On for Dogs [and Puppies Over 12 Weeks Old]]
[Sergeant's Evolve [11][21][40] Flea and Tick Squeeze-On for Dogs]
[Sentry®Pro XTR [11][21][40] Flea and Tick Squeeze-On for Dogs]
[Sentry®Pro XFT [11][21][40] Flea and Tick Squeeze-On for Dogs]
[Sentry®Pro XFTE [11][21][40] Flea and Tick Squeeze-On for Dogs]
[Sentry®Pro XFC+ [11][21][40] Flea and Tick Squeeze-On for Dogs]

[Optional marketing statements below may appear anywhere on the market label]

[Formulated for Dogs [and] Puppies over 12 weeks old [weighing] [9-20] [21-39] [40-60]] lbs.
[Flea & Tick Control for Dogs [&] Puppies 12 weeks old and older [weighing] [9-20][21-39][40-60]] lbs.
[For Dogs [&] Puppies Over 12 Weeks of Age [9 to 20] [21 to 39][40 to 60] (lbs)]
[[12][15][18][21][24][32][52] Week Supply Only For Use on Dogs and puppies [weighing] [9 to 20], [21 to 39], [40 to 60] (lbs)]
[Apply once every [4 weeks] [1 month] [30 days]!]
[Up to [4 week] [1 Month] Dose!]
[(ICON Showing up to 4 Week Protection)]
[Three Applications {for cartons with 3 tubes [vials]}] and/or [3 Month Supply] or [12 Week Supply]
[Four Applications {for cartons with 4 tubes [vials]}] and/or [4 Month Supply] or [16 Week Supply]
[Best if used year round!]
[Easy to Use Application]
[[Free] [brochure] [and] [Calendar] [Stickers] Inside!]
[Patent pending technology] [U.S. and Foreign patent pending technology]
[Dogs can be bathed 24 hours after squeeze-on is applied.]
[Quick Drying] [formula]
[with Vitamin E] [No oily residue]
[Fragrance Free]
[Formulated] [with] DERMASHIELD™ [Technology]
[DERMASHIELD™ [Technology] [which] dries quickly to keep the product on the outer skin surfaces]
[and contains [a vitamin] [vitamin E]]
[Not all dogs are the same. Should their flea and tick [control][product] be?]
[Brand Name] [developed for the weight of your dog] [formulated for the weight of your dog]

Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 9 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

[Multiple species]

[Kills fleas, ticks, flea eggs, flea larvae] [Repels mosquitoes]
[Kills fleas, ticks] [Kills flea eggs, flea larvae] [Repels mosquitoes]
[Four [4] Way Protection! Kills ticks, fleas, flea eggs & flea larvae]
[Five [5] Way Protection! Kills Fleas, Flea Egg, flea larvae, and ticks, and repels mosquitoes]
[Five in one protection! Kills fleas, flea eggs, flea larvae, and ticks, and repels mosquitoes]
[Kills all stages of the flea life cycle]
[Kills fleas, ticks, flea eggs, and flea larvae [and repels mosquitoes]]
[Kills fleas, flea eggs, flea larvae, ticks and repels mosquitoes]
[Kills fleas [in as little as 1 hour],[and] [ticks] [in as little as 3 hours], [flea eggs & flea larvae] [and repels mosquitoes]
[Kills Fleas [in as little as 1 hour],[and] Ticks [in as little as 3 hours]
[Kills Fleas, Ticks, Flea Eggs & Flea larvae [and repels Mosquitoes] for up to [4 Weeks] [1 month]]
[Works for up to 4 Weeks on fleas and ticks!]
[Protects for up to 4 Weeks on [adult] flea and tick]
[Up to 4 Week [One Month] [Adult] Flea and Tick [Control] [Protection!] [Treatment!]]
[Kills [Adult] Fleas and Ticks for up to 4 weeks!]
[Starts killing fleas and ticks in [less than] [just] [5] [15] minutes] [after application]
[Starts to kill fleas and ticks in [just] [5] [15] minutes] [after application]
[Stops][Prevents] fleas [from biting] [before they bite]
[Kills fleas before they bite]
[Kills fleas in [less than] [just] 1 hour [after application]
[Quickly kills flea in [just] [1 hour] and ticks [in as little as 3 hours]]
[Kills flea [as quickly as] [just] 1 hour [after application] and ticks [in as quick as] [just] 3 hours [after application]
[Kills Fleas, Ticks] [Kills Flea Eggs, Flea larvae] [Repels mosquitoes]

[Flea and Flea eggs and Larval]

[Starts killing fleas in [less than] [just] [5] [15] minutes] [after application]
[Starts to kill fleas in [less than] [just] [5] [15] minutes] [after application]
[Kills fleas in 1 hour] [after application]
[Stops][Prevents] fleas [from biting] [before they bite] [dogs][your dog]
[Stops][Prevents] fleas from ingesting a blood meal]
[Stops fleas from biting which may cause anemia]
[Stops fleas from biting [which may cause [flea allergy] dermatitis]
[[Stops][Kills] fleas which may cause flea allergy dermatitis]
[[Stops][Kills] fleas which may cause anemia]
[Kills fleas before they [can] bite]
[Quickly kills fleas in [less than] [just] [1 hour]]
[With Pyriproxyfen to Break Flea Life Cycle]
[With Insect Growth Regulator to Break Flea Life Cycle]
[With [Pyriproxyfen] Insect Growth Regulator which Breaks the Flea Life Cycle]
[Dual Action [!]: Cyphenothrin with Pyriproxyfen (or IGR), Effectively Breaks the Flea Life Cycle]
[Prevents Flea Eggs from Developing into Biting Adults]
[Breaks Flea Life Cycle!] for up to 9 weeks]
[Controls reinfestation for up to 9 weeks!]
[Kills Flea Eggs & Flea Larvae for up to [63 days], [9 weeks]!]
[Kills for up to [9 weeks] [63 days] [flea eggs] [flea larvae]]
[Kills fleas in as little as 1 hour!]
[Kills Fleas for up to 4 weeks!]
[Kills Adult Fleas for up to 4 weeks!]

Sergeant's Master Label

30 Oct 2017 Amendment
Information in [] is optional.

V17.1
Page 10 of 10

Reg. No. 2517-129
EPA SAL 05 Sep 2013

[Ticks]

[Starts killing ticks in [less than][just] [5][15] minutes [after application]

[Starts to kill ticks in [just] [5][15] minutes [after application]]

[Kills ticks in 3 hours] [after application]

[Quickly kills ticks in [as little as] [just] 3 hours][!]

[Kills and Detaches Ticks]

[Kills ticks in [as little as] [just] 3 hours][!]

[Prevents ticks from attaching and feeding [in as little as] [3 hours]]

[Kills ticks including those that may transmit Lyme disease]

[Kills Ticks for Up to [4 weeks][1 month][30 days]!]

[Kills Deer Ticks (vector of Lyme disease) for up to 4 weeks]

[Kills deer ticks [before they can] [that] [transmit Lyme disease] [for up to [4 weeks] [1 month][30 days]]]

[Kills Ticks (Including Deer Ticks) for up to [4 weeks][1 month][30 days]]

[Kills [brown dog ticks [(*Rhipicephalus sanguineus*)] [for up to [4 weeks][1 month][30 days]]]

[Kills American Dog Ticks [(*Dermacentor variabilis*)] for up to [4 weeks][1 month][30 days]!]

[Mosquitoes]

[Repels mosquitoes]

[Repels mosquitoes] [for up to [30 days] [4 weeks] [1 month]!]

[Repels mosquitos that may transmit heartworm]

[Repels mosquitos that may transmit [heartworm] [West Nile virus] [from feeding on dogs] [for up to [one month] [30 days] [4weeks]]]

[Repels mosquitoes] [Prevents mosquitos that may transmit West Nile virus and heartworm] [from feeding on dogs] [for up to [one month] [30 days] [4weeks]]]

[Prevents mosquitos from blood feeding, which may transmit viruses such as West Nile virus and heartworm to dogs for up to four weeks]