

WITH
WEED CONTROL
22-4-8

For ST. AUGUSTINE GRASS and
Certain Other Lawn Grasses

GUARANTEED ANALYSIS

| | |
|--|-----|
| Nitrogen (N) | 22% |
| Ammonium Nitrogen derived from Ammonium Sulfate | 1% |
| Water Soluble Nitrogen derived from Urea and other carbonyldehydes | 1% |
| Water Insoluble Nitrogen derived from Formaldehyde | 1% |
| Water Soluble Phosphate Acid (P_2O_5) | 4% |
| Derived from Phosphoric Acid | 4% |
| Soluble Potash (K_2O) | 4% |
| Derived from Monocal Phosphate | 4% |
| Total Available Plant Food | 34% |
| Chlorine, not more than | 0% |

Potential acidity equivalent to 1100 lbs. of Calcium Carbonate per ton.

No. 8-1014

Kills Weeds such as Chickweed, Dandelion, Knotweed, Plantain, Herbit, Spurge, Clover, Ground Ivy, Purslane, Ragweed, Morning Glory, Creeping Charlie and most other broadleaf weeds.

This product contains Trimec® Herbicidal Concentrate, a special formulation for use on St. Augustine grass. The patented mixture of herbicides gives maximum broadleaf weed control. This product can be used on sensitive grasses such as St. Augustine, bahia, centipede, bermuda, bent and zoysia. Under conditions of adequate moisture, high temperature and high humidity application may cause discoloration on St. Augustine. Under these conditions, however, the vigorous growth of St. Augustine will choke out most weeds and herbicide treatment would not be necessary. Do not apply to Dichondra, Lippia or similar ground covers.

When to apply. For best weed control, application should be made when weeds are growing vigorously. Application can be made at any time except for St. Augustine grass. Make application during spring and fall when temperatures are moderate.

How much to use. Apply evenly with a lawn spreader and do not care, mow or water the grass for at least 24 hours after spreading. This 25 lb. bag will cover 5,000 sq. ft. Close spreader openings when turning.

Grasses which are heavily infested by weeds may require more than one application. The product is suggested second application should be made in approximately 30 days.

CAUTION: Harmful if swallowed.

Avoid breathing or dust.
Avoid contact with skin, eyes, or clothing.
Wash thoroughly after use.
Do not store near fertilizer, seeds, insecticides or fungicides.
Do not use this product where delicate flowers are present or where cut flowers, vegetables or ornamental shrubbery.
Do not apply in areas underlain by roots of desirable trees and shrubs.
Do not contaminate irrigation, ditches or water used for irrigation purposes.
Do not reuse container. Destroy by burying in a safe place. Do not burn.

| CONTENTS | |
|------------------|------|
| Ammonium Sulfate | 6% |
| Boron | 0.5% |
| Chloride | 1% |
| Copper | 4% |
| Fertilizer | 0.5% |

PRINTED FOR
ACE HARDWARE CORPORATION

© 1980 ACE HARDWARE CORPORATION

Raoz
ACE