

ACCEPTED

FEB - 1 1989

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2548-28

MAX/KILL.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
(AND DOMESTIC ANIMALS)  
CAUTION**

Harmful if swallowed.  
Avoid inhalation spray mist.  
Irritating to skin and eyes.  
Do not get in the eyes.  
Avoid skin contact.  
Wash thoroughly with soap and water after using.  
Avoid contamination of cooking utensils, food ingredients, feed and water supplies.  
Do not use in the edible product areas of food processing plants.  
Do not use in milk houses or dairy barns.  
Do not spray directly on feed, feedstuffs or raw ingredients.  
Do not spray directly on pets, domestic livestock, especially dairy animals.  
Wear a pesticide respirator pump approved by the Mining Enforcement and Safety Administration formerly the U.S. Bureau of Mines and by the National Institute for Occupational Safety and Health under the provisions of CFR 20 Part II.

**ENVIRONMENTAL HAZARDS**

Remove fish, and birds and other domestic pets from room during spraying and keep out of room until spray is dry.  
This product is toxic to fish. Do not apply directly in water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**NON-WARRANTY**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

- PROMPTIONS**  
Do not contaminate water, food, or feed by storage and disposal.
- PESTICIDE DISPOSAL**  
Pesticide wastes are toxic. Improper disposal of excess pesticide or residue is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office for guidance.
- CONTAINER DISPOSAL**  
Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.
- GENERAL**  
Consult federal, state and local disposal authorities for approved procedures other than those given above. Approved procedures vary considerably for different types of generators.
- EQUIVALENT METHOD TO TRIPLE RINSING**  
Empty drums as completely as possible. Remove the lugs and place the drums upside down until liquid has evaporated.

**Mill & Bin Spray**

**Ready To Use - Long Lasting  
Quick Knockdown**

**KILLS EXPOSED STAGES OF:**  
Granary Weevil, Cadelle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Angoumois Grain Moth, Indian Meal Moth, Lesser Grain Borer, Roaches, Flies, Black Carpet Beetles & Meal Worms.

**ACTIVE INGREDIENTS:**  
Technical Methoxychlor\* ..... 3.0%  
Pyrethrins ..... .025%  
Technical Piperonyl Butoxide\*\* ..... 0.083%

**INERT INGREDIENTS:**  
Aromatic Petroleum Derivative Solvent ..... 20.0%  
Petroleum Distillate ..... 78.912%

\*Equivalent to 2.64% of 2, 2-bis (p-methoxyphenyl) -1, 1, 1-trichloro ethane and 0.36% of other isomers and related compounds.

\*\*Equivalent to 0.05% of (butyl carbityl) (4-propylpiperonyl) ether and 0.013% of related compounds.

**KEEP OUT OF THE REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** DO NOT INDUCE VOMITING. Seek medical aid. Aspiration of product into lungs due to vomiting can cause chemical pneumonitis which can be fatal. Emergency medical personnel should remove poison by gastric lavage.  
**IF BREATHED:** If inhaled in heavy concentrations or if illness results, remove to fresh air, and get medical attention. Administer artificial respiration, oxygen, or cardiopulmonary resuscitation if needed.  
**IF ON SKIN:** Remove contaminated clothing. Wash with mild soap and water. Launder contaminated clothing. If irritation occurs seek medical aid.  
**IF IN EYES:** Flush with large amounts of water for at least 15 minutes. If irritation persists, seek medical attention.

See side panel for precautionary statements.

EPA Reg. No. 2548-28

EPA Est. No. 2548-KS-1

NET CONTENTS

MANUFACTURED BY GALLONS



*Research Products Company*  
DIV. OF McSHARES, INC.  
SALINA, KANSAS 67401

11/87  
Scm to EPA

**DIRECTIONS FOR USE**

MAX KILL MILL & BIN SPRAY is a ready to use Combination long lasting insecticide that is best applied through conventional tank type pressure sprayers.

**AS A CLEAN UP SPRAY** in food processing plants, warehouses, mills and elevators apply as a coarse, wet spray to surfaces and areas where insects are known to be harboring. Avoid the contamination of equipment. After application allow for complete drying in areas where bagged feed or foodstuffs might be stored. It may be used as a clean up spray in bin bottoms after they have been thoroughly cleaned and before any grain is put back into the bin. One gallon should cover approximately 750 square feet.

**AS A PROTECTANT** in empty box cars, trucks and shipholds apply as a coarse wetting spray to all exposed surfaces. Allow for complete drying before any commodity loading.

**AS A BIN SPRAY** in farm granaries thoroughly clean and remove all dust and debris before applying a coarse, wetting spray to all exposed surfaces. Here again complete drying is necessary before any commodities are placed into the bin. Do not spray directly on grain and do not contaminate any feed or food stuffs. One gallon should cover approximately 750 square feet on non-absorbant surfaces.

**AS A FOG** in inedible product areas of food processing plants, elevator galleries and tunnels apply through a mechanical type fogger or mister at a rate of one quart per 5,000 cubic feet. Do not apply while equipment is in operation, and allow for complete settling of the fog or mist before any grain is conveyed. This method is quite satisfactory for the control of the adult stages of Indian Meal Moth. Repeat as needed.

See left panel for further precautions when using.