

WARNING

Keep Out of Reach of Children

HARMFUL IF SWALLOWED

Avoid inhalation spray mist.
Irritating to skin and eyes.
Do not get in the eyes.
Avoid skin contact.
Wash thoroughly with soap and water after using.
Avoid contamination of cooking utensils, food ingredients, feed and water supplies.
Do not use in the edible product areas of food processing plants.
Do not use in milk houses, dairy barns, or milk processing plants.
Do not apply inside bakery or food processing plants while in operation.
Cover food receptacles and utensils.
Do not spray directly on feed, foodstuffs or raw ingredients.
Destroy any contaminated food, feed or raw ingredients.
Do not spray directly on pets, domestic livestock, especially dairy animals.

ENVIRONMENTAL CAUTIONS

Remove fish, and birds and other domestic pets from room during spraying and keep out of room until spray is dry.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse containers. Destroy when empty. Keep containers closed when not in use. Keep away from heat and open flame. Wear approved safety equipment for protection of eyes, nose and mouth during application.

ANTIDOTE: Administer emetic of mustard. Call physician immediately.

Non Warranty

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

3% MALATHION

AIDS IN CONTROL OF:

Confused Flour Beetle, Red Rust Flour Beetle, Rice Weevil, Saw-Toothed Grain Beetle, Flat Grain Beetle, Lesser Grain Borer, Indian Meal Moth, Mediterranean Flour Moth, Angoumois Grain Moth, Granary Weevil, Cadelle Beetle, Roaches, Black Carpet Beetle, Silverfish, Ants.

ACTIVE INGREDIENTS:

Malathion (Prem. Gr.)* 3.0%
Pyrethrins05%
Technical Piperonyl
Butoxide ** 1.0%
Aromatic Petroleum
Derivative Solvent 10.0%
Petroleum Distillate 85.95%
100.00%

*O,O-dimethyl dithiophosphate
of diethyl mercaptosuccinate.

**Equivalent to .8% of (butycarbityl) (6-propylpiperonyl) Ether and .2% of related compounds.

EPA Reg. No. 2548-51

ACCEPTED

APR 17 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 2548-51

with comments

WARNING

Keep Out of Reach of Children

See additional warnings and antidote on side panels

CONTENTS

GALLONS

MANUFACTURED BY



Research Products Company

SALINA, KANS. 67401

DIRECTIONS FOR USE

FOOD HANDLING ESTABLISHMENTS - Places other than private residences in which exposed food is held, processed, prepared or served.

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY

Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facilities in operation also would be considered a food area).

Apply Max Kill 3% Malathion in small amounts directly into cracks and crevices using equipment capable of delivering a pin stream of insecticide in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases, in conduits, motor housings, junction or switch boxes where Confused Flour Beetle, Red Rust Flour Beetle, Rice Weevil, Saw Toothed Grain Beetle, Flat Grain Beetle, Lesser Grain Borer, Indian Meal Moth, Mediterranean Flour Moth, Angoumois Grain Moth, Granary Weevil, Cadelle Beetle, Roaches, Black Carpet Beetle, Silverfish and Ants hide. Prior to application de-energize electrical equipment. After application allow time for oil to evaporate before starting motors or placing switches in the "ON" position.

Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

NON-FOOD AREAS - Includes garbage rooms, lavatories, floor drains (to sewers, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage after canning and bottling).

Spot application with a coarse wet spray to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they may enter.

APPLICATION (General):

Apply Max Kill 3% Malathion with a low volume pressure tank type sprayer which gives a coarse wet spray with very little mist. Spray so as to thoroughly cover floors, walls, ceilings and all other surfaces where infestations may be present. Particular attention should be given to cracks and hiding or resting places. One gallon of Max Kill 3% Malathion should cover 750 square feet.

CEREAL GRAIN INSECTS: Max Kill 3% Malathion may be used for prevention and control of the exposed stages of cereal insects, weevils, various bran bugs and flour beetles, caddelles, moths, carpet beetles and meal worms. Apply as instructed under general application above.

INDIAN MEAL MOTH: Max Kill 3% Malathion may be used for prevention and control of Indian Meal Moth in bin tops and directly on the grain for moth control on grain surfaces of wheat, shelled corn, rye, barley, oats and grain sorghums. Not On Rice. Spray 1 gallon of Max Kill 3% Malathion per 750 sq. ft. in severe infestations. Break up webbing with a scissor before spraying. Spray adjacent bins and grain surfaces. Also spray galleys of the elevator or bin floors to control migrating moths and worms. Be sure surface grain is thoroughly mixed with rest of grain in bin when grain is removed from storage.

BROWN DOG TICKS IN PREMISES: Spray around baseboards, window and door frames, wall cracks and sleeping quarters of household pets. The frequency of application should depend on the source and the intensity of infestation. Fresh bedding should be placed in animal quarters following the spraying. Do not apply directly or indirectly to animals (See left panel).

CARPET BEETLES IN PREMISES: Spray localized areas of the floor and baseboards. Use oil can to squirt undiluted material into cracks and behind baseboards. Small areas under carpets may be sprayed but do not apply to large areas of the carpet. Spot application to floors, baseboards and shelves may be made and applications should be repeated as needed.

ANTS: Spray trails and areas around door sills and window frames where the pests may enter the premises. Spray openings around water pipes, heat ducts, electrical outlets and baseboards. Repeat applications as often as needed.

ROACHES: Apply thoroughly to infested cracks and other hiding or resting places. Spray a limited amount of exposed surface where roaches will crawl over it when they come out of hiding. Repeat applications as needed.

SILVERFISH: Apply as for roaches but in add treat baseboards, areas behind shelves, bookcases and range areas.