

PRECAUTIONARY STATEMENTS

Hazards To Humans (and Domestic Animals)

Irritating to skin and eyes.
Do not get in the eyes.
Avoid skin contact.
Wash thoroughly with soap and water after using.
Avoid inhalation of spray mist.
Do not remain in treated areas. Ventilate the area after treatment is completed.
Avoid contamination of cooking utensils, food, ingredients, feed and water supplies.
Remove fish and birds and other domestic pets from room during spraying and keep out of room until spray is dry.
Do not use in the edible product areas of meat packing or poultry processing plants.
Do not spray directly on pets, domestic livestock, especially dairy animals.
Do not give as internal treatment to animals.
Do not use in milk houses, dairy barns or milk processing plants.
Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.
Do not apply inside bakery or food processing plants while in operation.
Cover food receptacles and utensils.
Do not spray directly on feed, foodstuffs or raw ingredients.
Destroy any contaminated food, feed or raw ingredients.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Keep containers closed when not in use.
Keep away from heat and open flame.

ANTIDOTE:
External: Wash with soap and water.
Internal: Call physician immediately.

Storage and Disposal

PROHIBITIONS
Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
PESTICIDE DISPOSAL
Pesticide, spray mixture, or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
CONTAINER DISPOSAL
(a) Reseal container and offer for reconditioning, or
(b) Triple rinse (or equivalent) and:
offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.
GENERAL
CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES.

Non Warranty

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CONTACTICIDE 5

KILLS THE ACCESSIBLE STAGES ON CONTACT:

Granary Weevil, Rice Weevil, Cadelle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Angoumois Grain Moth, Indian Meal Moth, Cigarette Beetle, Drug, Store Beetle, Lesser Grain Borer, Flies, Cockroaches, Silverfish, Spiders, Ants, Crickets, Hornets, Wasps, Gnats, Mites, Mosquitoes.

ACTIVE INGREDIENTS. . . 100%
Pyrethrins. 0.5%
*Technical Piperonyl Butoxide. 1.25%
N-Octyl Bicycloheptene Dicarboximide. 2.25%
Petroleum Distillate. . . 96.00%
*Equivalent to 1.00% of (Butyl-carbityl) (6-Propylpiperonyl) Ether and .25% of related compounds.

EPA REG. NO. 2548-54
EPA EST. NO. 2548-K5-1

CAUTION

Keep Out Of The Reach Of Children

See side panels for additional Warnings and Antidote.

ACCEPTED

OCT 31 1979

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2548-54

NET CONTENTS

GALLONS

MANUFACTURED BY



Research Products Company
SALINA, KANSAS 67401

DIRECTIONS

MAX KILL CONTACTICIDE 5 is a surface and contact **READY-TO-USE** insecticide designed for use in flour mills, feed mills, rice mills, feed and food processing plants, cereal grain processing plants, grain elevators, granaries, shipholds, barges, empty boxcars, grain trucks, warehouses, feed and seed stores, grocery stores, restaurants and hotels.

WHEN USED AS A SPRAY in mills, cereal grain processing plants, grain handling facilities (elevators, granaries) for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, saw-toothed grain beetle, lesser grain borer, and other listed crawling insects, spray floors, walls and other surfaces of bins, storage and handling areas thoroughly, with particular attention to cracks, crevices, and similar protected locations. Waste grain, dust, dirt, and debris should be removed and destroyed, and storage areas thoroughly cleaned before spraying. Treat unloading, handling and processing areas, as well as **EMPTY** conveying, processing and handling equipment, for example the inside of reel sifters, purifier conveyors, sifters, roll housings and hoppers, pick-up conveyors, etc. Spray thoroughly around the base of heavy machinery and equipment. Let sprayed surface thoroughly dry before using. One gallon will cover 750 square feet of non-absorbent surface.

AS A PROTECTANT in empty boxcars, grain trucks, barges and shipholds for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, saw-toothed grain beetle, lesser grain borer and other listed crawling insects, spray the floors, walls, beams, cracks, crevices and other protected harborage points until surfaces are wet. A tanktype sprayer may be used utilizing low pressures. Allow for complete drying before loading. One gallon will cover 750 square feet of non-absorbent surface.

WHEN USED AS A SPRAY in food plants, grocery stores, hotels, feed and seed stores and restaurants, for the control of accessible stages of cockroaches, silverfish, spiders, ants, crickets, and listed cereal grain insects, use a sprayer which can operate at a low pressure and which will produce a coarse non-misting spray pattern. Avoid contamination of feed or foodstuffs as well as utensils or other food containers. Spray the floors, mold boards, cracks, crevices, and other areas where insects are known to hide. (See other precautions on side panel.) Apply spray until areas are wet but not to point of run-off. One gallon will cover 750 square feet of non-absorbent surface.

WHEN USED AS A FOG for indian meal moth, angoumois grain moth, hornets, wasps, gnats, mosquitoes, flies, and accessible stages of listed cereal grain insects in mills, elevators and empty granaries or grain bins, use a mechanical or air operated fogging apparatus and **MAX KILL CONTACTICIDE 5** at the rate of one quart per 5,000 cubic feet. Close all doors and windows before application and leave closed for 30 minutes after total dosage has been applied to allow for settling.

WHEN USED AS A FOG in empty boxcars, van type grain trucks, barges and shipholds for the control of listed flying insects and accessible stages of listed cereal grain insects, use a mechanical or air-operated fogging apparatus and **MAX KILL CONTACTICIDE 5** at the rate of one quart per 5,000 per cubic feet. Close all doors and hatches before applying and leave closed for 30 minutes after total dosage has been applied to allow for settling.

WHEN USED AS A FOG in boxcars or van type trucks containing bagged flour or feed for the control of listed flying insects and accessible stages of listed cereal grain insects, cover the bags with either paper or plastic tarps and apply at the rate of 8 ounces (1/2 pint) per 1,000 cubic feet of unoccupied space. Close all doors before applying and leave closed for 30 minutes after total dosage has been applied.

2548-54

10/31/1979

11