

ACCEPTED

2548-54

MAR 20 1985

Under the Federal Insecticide, and Fungicide, and Plant Disease Act

as a PESTICIDE and as a PRECAUTIONARY STATEMENTS registered under EPA Reg. No. _____

Hazards To Humans

(and Domestic Animals)

Irritating to skin and eyes
Do not get in the eyes
Avoid skin contact
Wash thoroughly with soap and water after using
Avoid inhalation of spray mist
Do not remain in treated areas. Ventilate the area after treatment is completed
Avoid contamination of casing utensils, food ingredients, food and water supplies
Remove fish and birds and other domestic pets from room during spraying and keep out of room until spray is dry
Do not spray directly on pets, domestic livestock, especially dairy animals
Do not give as internal treatment to animals
Do not use in milk houses or dairy barns
Do not apply directly to food
Do not apply space spraying while food processing is under way

With the exception of federally inspected meat and poultry plants, food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions indicated.

Destroy any contaminated food, food or raw ingredients

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Keep containers closed when not in use
Keep away from heat and open flame

ANTIDOTE

External: Wash with soap and water
Internal: Call physician immediately
If ingested: consult a physician. Do not induce vomiting unless directed by a physician. It can be hazardous to induce vomiting.

STORAGE AND DISPOSAL

- 1. PRECAUTIONS**
Do not contaminate water, food, or feed by storage or disposal.
- 2. PESTICIDE DISPOSAL**
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- 3. CONTAINER DISPOSAL**
Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
- 4. GENERAL**
Consult federal state and local disposal authorities for approved procedures other than those given above. Approved procedures vary considerably for different types of generators.
- 5. EQUIVALENT METHOD TO TRIPLE RINSING**
Empty drums as completely as possible. Remove the bung and place the drum upside down until liquid has evaporated.

Non Warranty

Some states no warranty expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

BEST AVAILABLE COPY

CONTACTICIDE 5

KILLS THE ACCESSIBLE STAGES ON CONTACT:

Granary Weevil, Rice Weevil, Cadelle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Angoumois Grain Moth, Indian Meal Moth, Cigarette Beetle, Drug Store Beetle, Lesser Grain Borer, Houseflies, Cockroaches, Silverfish, Spiders, Ants, Crickets, Hornets, Wasps, Gnats, Mites, Mosquitoes.

ACTIVE INGREDIENTS . . . 100%
Pyrethrins 0.50%
*Technical Piperonyl Butoxide 1.0
N-Octyl Bicycloheptene Dicarboximide 1.67%
Petroleum Distillate 96.83%
*Equivalent to 0.80% of (Butylcarbityl) (6-Propylpiperonyl) Ether and 0.20% of related compounds.

EPA REG. NO. 2548-34
EPA EST. NO. 2548-KS-1

CAUTION

Keep Out Of The Reach Of Children

See side panels for additional Warnings and Antidote.

NET CONTENTS

GALLONS

MANUFACTURED BY



RPC 2 '82

Research Products Company
SALINA, KANSAS 67401

DIRECTIONS

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTANT WITH ITS LABELING

MAX KILL CONTACTICIDE 5 is a surface and contact **READY-TO-USE** insecticide designed for use in flour mills, feed mills, rice mills, feed and food processing plants, cereal grain processing plants, grain elevators, granaries, shipyards, barges, empty silos, grain trucks, warehouses, feed and feed stores, grocery stores, restaurants and hotels.

WHEN USED AS A SPRAY in mills, cereal grain processing plants, grain handling facilities, elevators, granaries for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, saw-toothed grain beetle, lesser grain borer, and other listed crawling insects spray floors, walls and other surfaces of bins, storage and handling areas thoroughly, with particular attention to cracks, crevices, and similar protected locations. Waste grain, dust, dirt and debris should be removed and destroyed and storage areas thoroughly cleaned before spraying. Treat unloading, handling and processing areas, as well as EMPTY conveying, processing and handling equipment, for example the inside of rail cars, transfer chutes, elevators, hoppers, rail headings and hoppers, such as conveyors, etc. Spray thoroughly around the base of heavy machinery and equipment. Let sprayed surface thoroughly dry before using. One gallon will cover 750 square feet of non absorbent surface.

AS A PROTECTANT in empty boxes, grain trucks, barges and shipyards for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, saw-toothed grain beetle, lesser grain borer and other listed crawling insects spray floors, floors, walls, beams, cracks, crevices and other protected harborage points with surfaces are wet. A synthetic sprayer may be used utilizing low pressures. Allow for complete drying before loading. One gallon will cover 750 square feet of non absorbent surface.

WHEN USED AS A SPRAY in feed plants, grocery stores, hotels, feed and feed stores and restaurants for the control of accessible stages of contacticidal silverfish, spiders, ants, crickets, and listed cereal grain insects use a sprayer which can operate at a low pressure and which will produce a coarse non misting spray pattern. Avoid contamination of food or feedstuffs as well as animals or other food containers. Spray the floors, mold boards, cracks, crevices and other areas where insects are known to hide. Use other precautions on side panel. Apply spray until areas are wet but not to point of runoff. One gallon will cover 750 square feet of non absorbent surface.

WHEN USED AS A FOG for indoor use: moth, angoumois grain moth, hornets, wasps, gnats, mosquitoes, flies and accessible stages of listed insects in mills, feed processing plants, elevators and empty granaries or grain bins, use a mechanical or air operated fogging apparatus and **MAX KILL CONTACTICIDE 5** at the rate of one gallon per 5,000 cubic feet. Close all doors and windows before application and have closed for 30 minutes after total dosage has been applied to a satisfactory setting.

WHEN USED AS A FOG in empty boxes, grain trucks, barges and shipyards for the control of listed crawling insects and accessible stages of listed cereal grain insects use a mechanical or air operated fogging apparatus and **MAX KILL CONTACTICIDE 5** at the rate of one quart per 5,000 per cubic feet. Close all doors and windows before applying and have closed for 30 minutes after total dosage has been applied to allow for setting.

WHEN USED AS A FOG in places of other grain trucks containing bags of flour or feed for the control of listed crawling insects and accessible stages of listed cereal grain insects cover the bags with either paper or plastic tarp and spray at the rate of 8 ounces per pint per 1,000 cubic feet of unoccupied space. Close all doors before applying and have closed for 30 minutes after total dosage has been applied.