


ACCEPTED
 MAR 20 1985
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under
 EPA Reg. No. 2548-66

Case # 2548-66
 10/17/81
BEST AVAILABLE COPY



PRECAUTIONARY STATEMENTS
Hazards to Humans
(and Domestic Animals)
CAUTION

Irritating to skin and eyes
 Do not get in the eyes
 Avoid skin contact
 Wash thoroughly with soap and water after using
 Avoid inhalation of spray mist

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.
 If ingested, consult a physician. Do not induce vomiting unless directed by a physician. It can be hazardous to induce vomiting.

Do not remain in treated areas for more than 30 minutes after treatment is completed.

Avoid contamination of cooking utensils, food ingredients, food and water supplies.

Remove fish and birds and other domestic pets from room during spraying and keep out of room until spray is dry.

Do not spray directly on pets, domestic livestock, especially dairy animals.

Do not give as insect treatment to animals.

Do not use in fish houses or dairy barns.

Do not apply directly to food.

Do not apply space spraying while food processing is under way.

With the exception of federally inspected meat and poultry plants, food processing operations do not have to be stopped when applying a wet spray with care and in accordance with the directions and cautions indicated.

Destroy any contaminated food, food or raw ingredients.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as directed on the label.

PHYSICAL OR CHEMICAL HAZARDS

Keep container closed when not in use.
 Keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MAX HILL CONTACTICIDE 30 is a contact READY-TO-USE insecticide designed for use in flour mills, food mills, food and food processing plants, cereal grain processing plants, grain elevators, granaries, silos, bins, large, empty retailers' grain trucks, warehouses, food and food stores, grocery stores, restaurants and hotels.

CONTACTICIDE 30

KILLS THE ACCESSIBLE STAGES ON CONTACT:

Granary Weevil, Rice Weevil, Cadelle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Angoumois Grain Moth, Indian Meal Moth, Cigarette Beetle, Drug Store Beetle, Lesser Grain Borer, Flies, Cockroaches, Silverfish, Spiders, Ants, Crickets, Hornets, Wasps, Gnats, Mites, Mosquitoes.

ACTIVE INGREDIENTS
 Pyrethrins 0.30%
 Piperonyl Butoxide, technical 1.0%
 N-Octyl bicycloheptene dicarboximide 1.0%
 Petroleum distillates 97.7%
 *Equivalent to 0.8% of (Butylcarbityl) (6-Propylpiperonyl) ether and 0.2% of related compounds.

EPA REG. NO. 2548-66
 EPA EST. NO. 2548-KS-1

KEEP OUT OF THE REACH OF CHILDREN
CAUTION

See side panel for additional precautions and first aid.

NET CONTENTS

GALLONS

MANUFACTURED BY



Research Products Company

SALINA, KANSAS 67401

WHEN USED AS A SPRAY in mills, food processing plants, grain handling facilities, granaries, silos, bins, food and food processing plants, cereal grain processing plants, grain elevators, granaries, silos, bins, large, empty retailers' grain trucks, warehouses, food and food stores, grocery stores, restaurants and hotels, use a spray nozzle with a fan pattern. Spray the floors, walls, beams, cracks, crevices and other protected harborage points and surfaces to be treated. A fan-type sprayer may be used during low pressure. Allow for complete drying before loading. One gallon will cover 750 square feet of non-absorbent surface.

AS A PROTECTANT in empty bins, grain trucks, barges and shipholds for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, saw-toothed grain beetle, lesser grain borer and other listed pest insects, spray the floors, walls, beams, cracks, crevices and other protected harborage points and surfaces to be treated. A fan-type sprayer may be used during low pressure. Allow for complete drying before loading. One gallon will cover 750 square feet of non-absorbent surface.

WHEN USED AS A SPRAY in food plants, grocery stores, hotels, food and food stores and restaurants, for the control of accessible stages of cockroaches, silverfish, spiders, ants, crickets, and house flies, use a spray nozzle with a fan pattern at a low pressure and which will produce a coarse non-drifting spray pattern. Avoid contamination of food or foodstuffs as well as plants or other food components. Spray the floors, walls, beams, cracks, crevices, and other areas where insects are known to hide. Allow other procedures on side panel. Apply spray until areas are wet but not run to point of runoff. One gallon will cover 750 square feet of non-absorbent surface.

WHEN USED AS A FOG for Indian meal moth, angoumois grain moth, hornets, wasps, gnats, mosquitoes, flies and accessible stages of house flies in mills, food processing plants, elevators and empty granaries or grain bins, use a mechanical or air operated fogging apparatus and Max Hill Contacticide 30 at the rate of one quart per 1,000 cubic feet. Close all doors and windows before application and leave closed for 30 minutes after total dosage has been applied to allow for settling.

WHEN USED AS A FOG in empty bins, non-type grain trucks, barges and shipholds for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, lesser grain borer and lesser grain borer and lesser grain borer, use a mechanical or air operated fogging apparatus and Max Hill Contacticide 30 at the rate of one quart per 1,000 cubic feet. Close all doors and hatches before spraying and leave closed for 30 minutes after total dosage has been applied to allow for settling.

WHEN USED AS A FOG in bins or non-type trucks containing bagged flour or feed for the control of house flies, accessible stages of house flies and accessible stages of house flies, cover the bins with other paper or plastic tape and apply at the rate of 8 ounces (1/4 pint) per 1,000 cubic feet of unoccupied space. Close all doors before applying and leave closed for 30 minutes after total dosage has been applied.

STORAGE AND DISPOSAL

- 1. PRECAUTIONS**
Do not contaminate water, food, or feed by storage or disposal.
- 2. PESTICIDE DISPOSAL**
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- 3. CONTAINER DISPOSAL**
Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
- 4. GENERAL**
Consult federal, state and local official authorities for approved procedures other than those given above. Approved procedures vary considerably for different types of generators.
- 5. EQUIVALENT METHOD TO TRIPLE RINSING**
Empty drums as completely as possible. Remove the bung and place the drums upside down until liquid has evaporated.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than that indicated on the label. Buyer assumes all risk of use under handling of this material when such use under handling is contrary to label instructions.