

NEED DOCUMENT AVAILABLE

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(& DOMESTIC ANIMALS)
CAUTION**

Avoid eye and skin contact
Avoid dust inhalation
Wash thoroughly with soap and water after using
Wear approved respiratory equipment for protection of eyes, nose and mouth during application
Avoid contamination of feed and food

FIRST AID

If illness results from inhalation remove to fresh air and call a physician
If swallowed call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate natural water by cleaning of equipment or disposal of wastes.

NON WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE:
Keep containers closed when not in use. Do not use, pour, spill or store near heat or open flame.

DISPOSAL:

PESTICIDE DISPOSAL
Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

CONTAINER DISPOSAL
Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If container cannot be reused, dispose of in the same manner.

GENERAL
Consult Federal, state or local disposal authorities for approved alternative procedures.

MAX/KILL

DUSTACIDE 6

AID IN CONTROL OF:
Rice Weevil, Granary Weevil, Lesser Grain Borer, Saw Toothed Grain Beetle, Carlelle, Black Carpet Beetle, Flat Grain Beetle, Rusty Grain Beetle, Red Flour Beetle, Confused Flour Beetle, Mediterranean Flour Moth, Indian Meal Moth, Angoumous Grain Moth

ACTIVE INGREDIENTS:

Malathion	6.0%
(O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	
Inert Ingredients	94.0%
Total	100.0%

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See side panel for additional precautionary statements

**E.P.A. REG. NO. 2548-72
E.P.A. EST. NO. 2548-KS-1**

ACCEPTED
MAY 26 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2548-72



NET CONTENTS _____ **lbs.**
MANUFACTURED BY
Research Products Company
SALINA, KANSAS 67401

ACCEPTED
MAY 26 1982
EPA REG. NO. 2548-72

DIRECTIONS FOR USE

GENERAL

The primary function of Dusticide 6 is to protect stored grain from insect attack and to fumigate in order for the most effective protection must be uniformly mixed with the grain. The results achieved as the insects move about feeding after a surface application of Dusticide 6 will be effective against the adults of the common stored grain insects.

To control existing infestations it is best to fumigate the grain first and thereby kill the eggs, and then protect the surface against re-infestation with a surface application of Dusticide 6.

Dusticide 6 can be used as directed as a prophylactic treatment of the raw agricultural commodities - wheat, corn, sorghum, oats, barley, rice and rye.

DOSAGE

During transfer: 10 lbs. per 1000 bushel

Surface treatment: 5 lbs. per 1000 square feet

APPLICATION PROCEDURES

DURING TRANSFER. Calculate the quantity of Dusticide 6 required. Whether by hand or mechanical feeder apply Dusticide 6 uniformly and continuously as the grain is transferred. At the end of transfer protect the surface by applying 5 pounds per 1000 square feet.

SURFACE TREATMENT. Surface treatment is designed to protect uninfested grain from being infested and/or to control existing minor surface infestations. It is especially suited to protect the grain from attack by Indian Meal Moth. To be effective Dusticide 6 should be uniformly applied to the surface and raked in to a depth of approximately 6 inches.

NOTE: A well established Indian Meal Moth infestation will be difficult to bring under control short of fumigation. In either case it will be necessary to remove any webbing present and stir the surface of the grain before applying either Dusticide 6 or fumigating with a suitable fumigant. Any remaining webbing should be burned or buried.

DOCUMENT AVAILABLE