

Net Wt. 4 Lbs.

ZEE-TOX

SEVIN DUST 5%

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-Methylcarbamate)* 5.0%

INERT INGREDIENTS 95.0%

TOTAL — 100.0%

*Sevin Trademark of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate.

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Avoid contamination of feed or food stuffs.

DIRECTIONS:

Dust the upper and lower surfaces of the foliage thoroughly.

PEAS: Against Mexican bean beetle, leafhoppers, lygus, and root bugs, dust thoroughly, beginning when insects first appear and repeat every 7 to 10 days or as needed.

Apply 1 to 2 oz. per 50 foot row.

CUCUMBER, SUMMER SQUASH: Against striped and spotted cucumber beetle, squash bugs, flea beetle, pickleworms and other foliage pests. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

TOMATO, EGGPLANT, PEPPER: Against tomato fruitworm, fall armyworm, and flea beetle, dust thoroughly when insects first appear and every 7 to 10 days if necessary. Use 1 to 2 oz. per 50 foot row.

POTATO: Against Colorado potato beetle, leafhopper, and flea beetle apply 1 to 2 oz. per 50 foot row.

COLLARDS, TURNIPS (tops), MUSTARD GREENS: Against flea beetle, corn earworm, Armyworms, Imported cabbage worm use 1 to 2 oz. per 50 foot row when insects first appear and repeat every 7 to 10 days if necessary for control.

DOGS AND CATS: For fleas and ticks apply dust liberally over entire animal, avoiding eyes, and rub thoroughly into skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if necessary. Avoid treatment of kittens less than 4 weeks of age.

Dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or powder duster. Force dust into cracks and crevices. Repeat as necessary.

Do not use this product on poultry. Do not reuse container. Destroy when empty.

This product is highly toxic to bees when exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agriculture Extension Service.

NOTE TO PHYSICIANS: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Packed By

**BUHL CHEMICAL
ORLANDO, FLORIDA**

EPA EST. NO. 2553-FL-1

EPA REG. NO. 2553-33

