


HARTZ FLEA & TICK KILLER

Contains Allethrin

FOR CATS AND DOGS

- kills fleas, ticks and lice
- stops scratching
- repels gnats, mosquitoes and flies
- deodorizes



Net Wt. 8 oz 226 g

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep this and all other containers of this product out of the reach of children. Avoid contact with eyes, nose, mouth, and skin. Wash hands thoroughly after use. Do not use in areas where food is prepared or stored.

PHYSICAL OR CHEMICAL HAZARDS

This product is not flammable. It is not a strong irritant. It is not a strong oxidizer. It is not a strong reducer. It is not a strong acid. It is not a strong base.

DIRECTIONS FOR USE

Shake well before use. Spray over the animal, beginning at the tail and moving toward the head. On long-haired dogs and cats, ruffle the coat and spray against the lay of the hair. Spray directly to the skin so that spray reaches skin. For best results in controlling tick infestations, locate ticks and spray directly on them. Avoid spraying face and genitals. An application of 30 to 45 seconds is usually sufficient for an average size dog; for cats about 10 seconds is adequate. Repeat application weekly if necessary. It is recommended that the pet's bedding or sleeping quarters and areas and surfaces in which the pet is likely to be treated for complete flea and tick control and to keep the pet's quarters.

STORAGE AND DISPOSAL

Do not incinerate. Wrap container and put in trash. Do not reuse.

Remove protective cap, hold container upright, move away from you. With nozzle 10 to 12 inches away, direct spray over the animal, beginning at the tail. Spray over the animal until the entire coat is damp, up to the back of neck. On long-haired dogs and cats, ruffle the coat and spray against the lay of the hair. Spray directly to the skin so that spray reaches skin. For best results in controlling tick infestations, locate ticks and spray directly on them. Avoid spraying face and genitals. An application of 30 to 45 seconds is usually sufficient for an average size dog; for cats about 10 seconds is adequate. Repeat application weekly if necessary. It is recommended that the pet's bedding or sleeping quarters and areas and surfaces in which the pet is likely to be treated for complete flea and tick control and to keep the pet's quarters.

ACCEPTED

SEP 15 1980

2596-10

17