

[Note: Optional text in parentheses or brackets]

FRONT PANEL

**Hartz 2 in 1 Flea & Tick Collar For Dogs**

Alternate Trade Names:

Hartz 2 in 1 Flea & Tick Collar For Large Dogs  
Hartz 2 in 1 Flea & Tick Collar For Puppies

[Bright (Purple) (Red) (Blue) (White)]

(with Rabon)  
(with Deodorant)  
(Long term protection)

(For Dogs)  
(For Puppies)  
(For Large Dogs)

(Kills for 5 Months)

CAUTION: Do not let children play with this collar.  
See back panel for ingredient statement, precautionary statement, and directions for use.

CONTENTS: 1 (purple) (red) (blue) (white) collar, Net Wt. \_\_\_\_ oz. \_\_\_\_ g

SIDE, TOP, or BOTTOM PANELS


Hartz 2 in 1 Flea & Tick Collar [for (large) dogs (puppies) with necks to (15") 20" (23") (26")]

(purple) (red) (blue) white collar

To activate collar, hold at each end ...STRETCH...it's ready to work

Powder action

Unconditional money back guarantee by Hartz



BACK PANEL

Kills fleas and ticks common to dogs, including Rocky Mountain Wood Tick, carrier of Rock Mountain Spotted Fever and the Deer Tick, which may carry Lyme Disease

ACTIVE INGREDIENT:	BY WEIGHT (Nom.)
tetrachlorvinphos (CAS # 22248-79-9) .....	14.55%
OTHER INGREDIENTS .....	85.45%
TOTAL.....	100.00%

READ ENTIRE LABEL BEFORE EACH USE  
USE ONLY ON DOGS

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID – Contains an organophosphate that inhibits cholinesterase. IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow . Do not induce vomiting unless told to do so by a Poison Control Center or doctor Do not give anything by mouth to an unconscious persons. IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice. IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

Note to Physicians and Veterinarians: This product contains an organophosphate and may cause cholinesterase inhibition. Cholinergic symptoms may include salivation, miosis, incoordination, muscle fasciculation and/or weakness, vomiting, and diarrhea. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine. Call 1-800-275-1414 or your local Poison Control Center for further information.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**REMOVE COLLAR FROM PACKAGE, UNROLL AND STRETCH TO ACTIVATE INSECTICIDE GENERATOR.**

Do not use on puppies under 6 weeks of age. Do not unroll collar until ready to use. Place the Hartz 2 in 1 Flea & Tick Collar around the dog's neck, adjust to proper fit, and buckle in place. The collar must be worn loosely to allow for growth of the dog and to permit the collar to move about the neck. Generally, a properly fitted collar is one that when fastened will snugly slide over the dog's head. Leave 2 or 3 inches on the collar for extra adjustment and cut off and dispose of extra length.

Consult a veterinarian before using this product on debilitated, aged, or medicated animals. If used on pregnant or nursing dogs, do not replace collar until the puppies are at least 6 weeks old. Sensitivities may occur after using ANY pesticide product for pets. Some animals may become irritated by any collar if it is applied too tightly. If this occurs, loosen the collar. If irritation continues, remove the collar from the animal. If signs of sensitivity occur, remove the collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea & tick collars may be immediately replaced.

This collar is intended for use only as an insecticide generator. The collar will begin to kill fleas and ticks immediately. Its full protection against harder to kill ticks will be built up within a few days after being placed on the pet. For continuous flea and tick protection under normal conditions replace the collar every 5 months. Under conditions where dogs are exposed to severe flea or tick infestations it may be necessary to replace more frequently, however, do not replace the collar more frequently than every 4 months. Wetting will not impair the collar's effectiveness or the pet's protection. If the dog goes swimming or is out in the rain it is not necessary to remove the collar. The Hartz 2 in 1 Flea & Tick Collar may be worn with a regular collar.

**STORAGE and DISPOSAL**

Storage: Store at room temperature.

Disposal: Do not use empty pouch. Place in trash or offer for recycling if available.

For more information call our flea and tick experts at 1-800-275-1414 weekdays, 9 am – 5 pm E.S.T.

EPA Reg. No. 2596-62

EPA Est. No. 2596-\_\_\_\_\_

(Important: Consumer information and pet I.D. tag offer printed on inside of package.)

Visit our website [www.hartz.com](http://www.hartz.com)

Hartz Every Pet's Best Friend

Made in the U.S.A. by  
©The Hartz Mountain Corporation, Secaucus, NJ 07094