

**ACCEPTED**

NOV 20 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2596-79

**HARTZ  
2 in 1  
FLEA AND TICK  
POWDER FOR DOGS**

Alternate Trade Name:

**HARTZ® 2 in 1 ADVANCED CARE™ FLEA AND TICK  
POWDER FOR DOGS**

**USE ONLY ON DOGS.**

**ACTIVE INGREDIENT:**

(Z)-2-chloro-1-(2,4, 5-trichlorophenyl) vinyl dimethyl phosphate  
(Tetrachlorvinphos) (CAS # 22248-79-9):.....3.3%  
OTHER INGREDIENTS:.....96.7%

**Keep out of reach of children.**

**CAUTION:** See back panel for precautionary statements.

**FIRST AID** – Contains an organophosphate that inhibits cholinesterase.

**IF SWALLOWED:** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

**Note to Physicians and Veterinarians:** This product contains an organophosphate and may cause cholinesterase inhibition. Cholinergic symptoms may include salivation, miosis, incoordination, muscle

fasciculation and/or weakness, vomiting, and diarrhea. Atropine is antidotal only if symptoms of cholinesterase inhibition are present. 2-PAM may be effective as an adjunct to atropine. Call 1-800-275-1414 or your local Poison Control Center for further information.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling. Wash hands before eating, chewing gum, using tobacco or using the toilet. Not for use on puppies less than 12 weeks of age. Consult a veterinarian before using this product on debilitated, aged or medicated animals. Sensitivities may occur after using ANY pesticide product on pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea and tick collars may be immediately replaced.

**BACK PANEL**  
Mandatory Copy

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL BEFORE EACH USE.**

To kill fleas, ticks and lice, and to reduce itching and scratching due to insect bites, dust entire dog beginning at head and working back. Make sure that the powder gets down to the skin. Take care to treat feet and legs.

**Recommended Dosage:** Use approximately 1/2 ounce of powder for a small dog (e.g. Terrier). Use approximately 1 ounce of powder for a medium dog (e.g. Spaniel). Use approximately 1 1/4 ounce of powder for a large dog (e.g. Labrador). More powder may be needed for longhaired dogs. There is sufficient powder in this container to treat a small dog 4 times.

2596-79  
06/25/03

2596-79

11/20/2003

1/2

Lightly dust dog's bedding with approximately the same amount of powder. Repeat treatment of dog and bedding at weekly intervals if necessary.

To control Sarcoptic Mange – a problem often encountered by dogs. Lesions which form are usually characteristic by loss of hair, redness and possible open sores. Hartz 2 in 1 Flea and Tick Powder will kill the mite which causes this problem. To treat the dog, apply the powder as directed above, making sure that all sores and lesions are covered. Repeat application weekly for at least 3 treatments. More advanced lesions may require further weekly treatments. The area will begin to heal immediately, but complete healing, including hair growth, may take several weeks. If irritation persists, consult your veterinarian.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage and disposal. Store in a cool, dry place. **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Storage: Store at room temperature.

EPA Reg. No. 2596-79  
EPA Est. No. 2596-NJ-1

Net Wt. 4oz (113 g)

The Hartz Mountain Corporation  
Secaucus, NJ 07094 and St. Thomas, Ontario N5P 3W7 Canada

Made the USA

\*Rabon is the Hartz trademark for Tetrachlorvinfos  
Visit our website: [www.hartz.com](http://www.hartz.com)

**Optional Copy: Front or Back Panel**

DIRECT TREATMENT  
With Rabon\*  
Kills Fleas, Kills Ticks

Unaffected by Water  
Kills Fleas and ticks, including the deer tick, which may carry LYME DISEASE

KILLS LICE WHEN A PROBLEM \* FAST ACTING \* LASTS 7 DAYS \*  
UNAFFECTED BY WATER \* KILLS INSECTS WHICH MAY CAUSE  
ITCHING AND SCRATCHING

CONTROLS SARCOPTIC MANGE

Hartz - Every Pet's Best Friend  
For more information, call our flea and tick experts at 1-800-272-1414  
weekdays, 9AM - 5 PM EST.