

REPELS FLIES, GNATS, AND MOSQUITOES
KILLS FLEAS, LICE, AND TICKS

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION: Do not use near birds, fish or foodstuffs. Do not spray in eyes or face of cat. Do not use on kittens under 6 weeks old. Wash hands thoroughly after use.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. Hold bottle upright 6 inches from pet. Direct spray toward pet and spray entire coat, pressing dispenser with quick, short strokes. Move bottle to get even coverage of coat (until tips of hair are moist).
2. Apply lightly and rub into animal's coat. Ruffle hair for spray to reach skin. Spray 4 to 6 seconds to a five pound cat, vary according to weight of cat.
3. Dry with towel.
4. Comb and brush coat.
5. Repeat once a week if necessary.
6. Kills fleas in 5 to 10 minutes. Ticks are tough--spray directly.

Storage and Disposal

Do not re-use empty container. Wrap container and put in trash collection.

EPA Reg. No. 2596-93

EPA Est. 34961-NJ-01

Distributed by

© THE HARTZ MOUNTAIN CORPORATION

Harrison, N.J. 07029-9987 and St. Thomas, Ontario N5P 3W7 Canada

Made and Printed in U.S.A.

HARTZ

**CAT
FLEA & TICK
KILLER**

DEODORANT
REPELLENT

FINE MIST SPRAY

EPA Reg. No. 2596-93
EPA Est. 11525-NJ-1

ACTIVE INGREDIENTS	
Pyrethrins	.060%
Piperonyl Butoxide-Technical*	.120%
N-Octyl bicycloheptene	200%
Baccharoside	288%
Petroleum Distillate	99.332%
INERT INGREDIENTS	100.000%

*Equivalent to 0.096% (butylcarbitol)
(5 Propyl-piperonyl ether and 0.024%
related compounds)

Contains Sodium Nitrate

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements

8 fl oz 236 ml

2596-93

CROP	DISEASE	POUNDS TERRACLOR® 75% W.P.	REMARKS
Magnolia grandiflora	Magnolia Leaf Spot (Phyllosticta cockeri)	2	Apply specified dosage in 100 gallons of water as a broadcast foliar spray. Add a spreader-sticker at the rate of 1 pint per 100 gallons. Begin application approximately 1 week before disease normally appears. Apply at least 4 sprays at 2 week intervals. Note: Do not use on Magnolia fuscata as injury may result.
Roses	Botrytis Storage Rot	1 to 2	Dip Application: Mix specified dosage in 100 gallons of water and dip dormant roses prior to storage. Maintain good agitation in dip tank.
		2	Spray Application: Mix specified dosage in 100 gallons of water thoroughly spray dormant roses prior to storage.
Southern Pine Seedlings	Rhizoctonia Needle Blight	50	Apply specified dosage per acre as a broadcast spray in 100 gallons of water. An additional 1/2 inch of water per acre should be applied by irrigation. Seed may be planted immediately after irrigation or within one week of this date.
Lawns and Turf	Brown Patch (Rhizoctonia solani)	See Remarks	For Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses: Make one application in the fall or spring at first indication of infection. Mix 1 lb. in sufficient water (10-15 gals.) to cover 1000 sq. ft. Apply by means of a sprinkler can, sprayer or any other convenient method of obtaining even coverage. Following application, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3-4 weeks later. For Dichondra: Treat as above, using 1 lb. mixed in 40 gals. of water per 1000 sq. ft.