

This product fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance 1965 Recommendations of the U. S. Public Health Service in waters up to 500 p.p.m. of hardness, calculated as CaCO<sub>3</sub> when tested by the A.O.A.C. Germicidal and Detergent Sanitizers Official Method.

**ACTIVE INGREDIENTS**

|   |               |
|---|---------------|
| n-Alkyl (C <sub>14</sub> , 95%, C <sub>12</sub> 3%, C <sub>16</sub> , 2%) dimethyl benzyl ammonium chloride dihydrate | 50.0%         |
| Ethylene diamine tetraacetic acid   | 9.0%          |
| <b>INERT INGREDIENTS</b>  | <b>41.0%</b>  |
|   | <b>100.0%</b> |

**Sanitabs**  
**SANITIZING TABLETS**  
**100 TABLETS**

**DANGER**

**DIRECTIONS**

Do not use with soap or synthetic detergent. Avoid contamination of food. Scrape utensils and empty glasses before washing. For Sanitizing Eating and Cooking utensils in Restaurants, Soda Fountains, Taverns, etc.

1. Utensils must be thoroughly washed in a hot detergent solution which is kept reasonably clean. (First compartment)
2. Utensils must be then rinsed free of such solution. (Second compartment)
3. Utensils must be then immersed in a clean sanitizing solution for at least 1-2 minutes. This solution is to be maintained at the proper concentration at least 200 ppm quaternary and minimum temperature of 75 F. (water hardness not to exceed 500 ppm at pH of 5.0 or higher. (Third compartment)

Allow to air dry. Do not use. Two (2) Sanitabs per 3 gallons of water make a 200 ppm solution. Sanitabs should be diluted in warm or hot water. Stir solutions well before using. EPA Reg. No. 2626-25

**ACCEPTED**

July 10 1974

UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 2626-25

B. Q.  
*Sanitabs*  
SANITIZING  
TABLETS

Net Weight 8 oz

**DANGER**

KEEP OUT OF THE REACH  
OF CHILDREN

SEE RIGHT (SIDE) PANEL FOR  
FIRST AID AND ADDITIONAL  
PRECAUTIONS



DIRECTIONS

ACCEPTED

CHEMICAL RESEARCH PRODUCTS  
NORFOLK VIRGINIA