

[2666-38]

Pyrenone "Fog" Spray is accepted for use in federally inspected meat and poultry processing establishments.

CAUTION: Do not spray into open flame, or on green plants. Harmful if swallowed. Keep from reach of children. To avoid any explosion hazards, not more than 1 gallon of oil should be atomized in each 40,000 cubic feet, pilot lights should usually be extinguished before fogging, and not relighted until room is aired.

NOTICE: Since weather and other conditions under which this product may be used will vary, Seller makes no warranty of any kind, express or implied, concerning use of this product. The user assumes all risks of use or handling when not used in accordance with directions.

PYRENONE "FOG" SPRAY kills on contact. For best results, use in a good fog generator (such as the Dyna-Fog). Treat the infested areas thoroughly and repeat application as needed.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.



U.S.D.A. Reg. No. 2666-38

PYRENONE* "FOG" SPRAY

ACCEPTED
Aug-31-1971
UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 2666-38 SUBJECT TO ATTACHED COMMENTS. 38

CAUTION:

KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

*Registered trademark of U.S. Industrial Chemical, Inc.

ACTIVE INGREDIENTS:	100.00%
**Technical Piperonyl Butoxide	5.03%
Pyrethrins	.62%
Petroleum Distillate	94.35%

**Equivalent to 4.02% of butylparbityl 6-propyl-piperonyl ether and 1.01% of related compounds.

NET CONTENTS

CORNELL PYRENONE* "FOG" SPRAY kills accessible stages of Flour Beetles, Rice Weevils, Cadelles, Saw-Toothed Grain Beetles, Mediterranean Flour Moth, Indian Meal Moths, Angoumois Grain Moth, Fried Fruit Beetles, and Grain Mites affecting stored grain and food products, and kills Cock roaches, Ants, and Silverfish in storage warehouses and food handling establishments. The proper use of this spray is an excellent adjunct to spot fumigation operations.

To obtain the best results in controlling Flying Insects, such as flies, gnats, mosquitoes, close doors and windows, create a dense fog in the room and leave the room closed for at least 15 minutes. Food receptacles, kettles and mixing equipment should be covered before the room is fogged to keep the dead insects from falling into the equipment.

For Crawling Insects, a heavier fog is necessary to penetrate where the insects may be hiding. It is advisable to thoroughly clean all areas to be sprayed. Avoid direct spraying on any food products.

This concentrate contains Piperonyl Butoxide and Pyrethrins. Applied according to directions, it can be used around food and food products, food handling utensils and equipment, and storage areas for food and feeds. Exposed food should always be covered before fogging operations start.

Do not remain in treated areas and ventilate the area after treatment is completed.

CORNELL CHEMICAL & EQUIPMENT CO., INC.
1115 NORTH ROLLING ROAD
BALTIMORE, 21228 MARYLAND, U.S.A.