Front Panel

MEM

DAIRY AND LIVESTOCK DUST

Contains: malathion and methoxychlor

ACTIVE INGREDIENTS: Methoxychlor, technical*	5.0%
Malathion O, O-dimethyl phosphorodithioate of diethyl mercaptosuccinate	4.0%
*Equivalent to 4.4% 2,2-bis(p-methoxyphenyl)-1,1,1-	
trichloroethane and 0.6% related compounds.	

Keep Out Of Reach Of Children

CAUTION

See Back Panel For Additional Cautions

NET CONTENTS: 1 1b. (453.6 g)

ACCEPTED

JUN 22 1989

Under the Federal Inserticide, Fungicide, and Rodenticide Act, as amended, for the posticide regimered under EPA Reg. No. 2724 -26

M&M is a trademark of Sandoz Ltd.

	····

	•••••
•••••	******
	•
	•••••

*****	•
•••••	•
	. • • . • .
	•
	•:•••
	·•• •

Back Panel

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS - CAUTION: Avoid contact with skin, eyes, and clothing. Avoid breathing dust. Wash thoroughly after using. Do not apply within five hours before or during milking.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e. swamps, bogs, marshes and potholes). Drift and runoff may be hazardous to aquatic organisms near the application site.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For direct application to Dairy Cattle and Beef Cattle, for control of Horn Flies: Apply 2 heaping tablespoonsful (2 oz.) to the backs and necks of infested animals. Rub lightly into hair. Repeat application in 10 to 14 days if necessary. On dairy animals, repeat no more often than once in every three week period, if necessary.

For control of Lice and Ticks on Beef Cattle: Treat entire animal very thoroughly. Repeat treatment at two week intervals if necessary. May be used on non-lactating dairy animals for lice control up to within two weeks of freshening.

As an aid for the control of Face Flies on Beef Cattle: Dust the faces of cattle thoroughly. Frequent repeated applications may be necessary.

For control of Hog Lice: Apply to hogs, pens, and litter. One treatment may be sufficient. Treat animals thoroughly. Repeat in 10 days or thereafter as needed.

<u>Lice and Keds on Sheep</u>: Treat animals thoroughly, preferably following shearing. Repeat as necessary.

STORAGE & DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of the material when such use and handling are contrary to label instructions.

Zoecon Corporation
A Sandoz Company
12200 Denton Drive, Dallas, Texas 75234

EPA Reg. No. 2724-76

1188-B: 6443A

(

Made in USA 1984 - Loecon EPA Est