


1(072)



# LINTOX-M

(CONTAINS MALATHION AND TOXAPHENE)

## LIVESTOCK SPRAY AND DIP

AND BACKRUBBER REFILL



QUICK KILL / LONG ACTING

ONE GALLON MAKES UP TO 200 GALLONS OF SPRAY

ACTIVE INGREDIENTS	
TOXAPHENE (CHEMICAL ANALOGUOUSLY GAMPHRENE)	42.0%
MALATHION (D-D METHYLENE DIPHOSPHORIC ACID DIMETHYL ESTER)	43.0%
PETROLEUM SULFONATE	39.8%
INERT INGREDIENTS	12.9%
	100.0%

**WARNING:** Keep Out of Reach of Children

See back panel for additional warnings

NET CONTENTS: FIVE GALLONS

EPA REG. NO. 272454-AA-11767

### INSECTS CONTROLLED

- Horn Fly
- Stable Fly
- Lone Star Tick
- Gulf Coast Tick
- Cattle Fever Tick
- Winter Tick
- Ear Tick
- Sheep Tick
- Sheep Scabies

Manufactured by Thuron Industries, Inc.

12200 DEHTON DRIVE, DALLAS, TEXAS 75234

2(0f2)

ACCEPTED  
4-2-74  
2724-54



TAKE TIME

OBSERVE LABEL  
DIRECTIONS

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROENTGENIC  
ACT FOR ECONOMIC POLLUTION REGULATION  
ED UNDER 20.724-54

# DIRECTIONS FOR USE

With Cermine

**ON BEEF CATTLE AND HORSES** - For control of Horn Fly, Cattle Lice, Ear Tick, Horn Fly, Tick, Gulf Coast Tick, Winter Tick, Cattle Fever Tick, mix one gallon of LINTOX-M with 100 gallons of water and shake well to form a creamy emulsion. Spray animals with a strong spray to penetrate to the skin or dip thoroughly. Approximately two quarts of spray are required for all animals and proper treatment rates for younger animals. Be sure not to overdose young animals.

For maximum long-lasting protection, mix one gallon of LINTOX-M with 100 gallons of water as described above. However, for a lower cost spray for temporary control of Cattle Lice and Horn Flies, one gallon of LINTOX-M may be mixed with 150 or 200 gallons of water and applied as above.

**ON SHEEP AND GOATS** - For control of Goat Lice, Sheep Lice, Keds, mix one gallon of LINTOX-M with 200 to 400 gallons of water, and spray or dip animals thoroughly. Be sure to wet or moisten thoroughly.

**ON HOGS** - For control of Hog Lice and Sarcopic Mange, dilute one gallon of LINTOX-M with 100 gallons of water, mix well, and spray or wash animals thoroughly. Be sure not to overdose young animals.

**MANGE AND SCAB** - For scab and prevention of Sarcopic, Psoroptic, and Chorioptic Mange (scab) on cattle, sheep and hogs, mix one gallon of LINTOX-M with 100 gallons of water and dip animals thoroughly. One thorough treatment is usually sufficient. Apply second treatment in 10 days if needed.

**ON BACKRUBBERS** - For use on burins, sark or chain backrubbers as an aid to controlling Horn Flies and Lice on beef cattle, dilute one gallon of LINTOX-M with 10 to 20 gallons of fuel oil or other suitable carrier and fill insecticide compartment of backrubber device or soak sack or cloth as the operator requires. Recharge as necessary. Rinse backrubber device when animals land water, travel regularly or trail. Do not allow beef cattle access to backrubber within 28 days of slaughter.

As a premise spray for barnyard and buildings - to kill Fleas, Mosquitoes, Roaches, Stable Flies, around barns, barnyards and outbuildings - dilute one gallon of LINTOX-M with 20 gallons of water. Thoroughly wet all surfaces where insects alight or crawl. For Fleas, thoroughly treat places where animal's sleep. Repeat applications as necessary for effective season-long control. Do not treat pets or animals with this mixture. Do not use in dwellings.

**CAUTION** - As a general precaution in spraying animals, avoid treating during cold, stormy weather, avoid treating overheated or sick animals; water animals before dipping to prevent them from drinking the dip. If calves must

be treated, spray with a fine mist, not a stream.

**WARNING** - May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or clothing. Wash thoroughly. Do not apply in dairy barns or to dairy animals. Avoid contamination of feed and feedstuffs. Do not use on dogs, cats or other household pets, or around fish. Do not use on animals within 28 days of slaughter. Do not treat animals under three months of age. If young animals are to be dipped, only just dipping is recommended to avoid the possible swallowing of the dipping solution by the animal. Do not treat sows within 2 weeks of farrowing or for at least 3 weeks thereafter. Do not use in poultry houses or on poultry. Do not reuse container. Destroy when empty.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Melphar is a cholinesterase inhibitor. Do not use this product simultaneously with other drugs, insecticides, pesticides or chemicals having cholinesterase inhibiting activity nor within a few days before or after treatment with any other cholinesterase inhibitor.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label instructions.

## LINTOX-M DILUTION TABLE

Amount of LINTOX-M	Gallons of Water				
	Cattle, Horses, Hogs	Sheep and Goat Lice and Keds		Sheep Scabies	Premise Spray
		Spray	Dip		
Quart	25 Gal.	50 Gal.	100 Gal.	75 Gal.	5 Gal.
Gallon	100 Gal.	200 Gal.	400 Gal.	100 Gal.	20 Gal.

THURON INDUSTRIES, INC. 1957

Made in USA

## INSECTS CONTROLLED

Cattle Lice

Sheep Lice

Goat Lice

Hog Lice

Hog Mange

Cattle Mange

and

Many Other

Insect Pests

Manufactured by Thuron Industries, Inc.

12200 DENTON DRIVE, DALLAS, TEXAS 75234

ACL 00460-D