

BEEF CATTLE: To control Lice, Ticks and Hor Flies use at the rate of 2 quarts per 50 gallons of water. Apply as a complete coverage spray. Repeat every two weeks if necessary.

POULTRY: To control Northern Fowl Mite, Poultry Lice and as a supplement to premise treatments for Chicken Red Mite use 2 tablespoons per 1 gallon of water. Spray 1 gallon of dilute mixture per 100-150 birds.

POULTRY HOUSES: To control Northern Fowl Mite, Chicken Red Mite, Poultry Lice, Flies and Mosquitoes, use 4 tablespoons per 1 gallon of water. Apply to walls, ceilings, roost, nests, and cracks and crevices. Repeat as necessary.

FLIES AND MOSQUITOES: Use at the rate of 2 quarts per 25 gallons (1 pint to 6 gallons) of water. For best results add 1 pound of sugar for each gallon of finished spray. Mix well. Spray all surfaces and places where Flies and Mosquitoes light or congregate. Use a wetting spray and spray to the point of run off. Use around buildings, stockyards, dairy barns, poultry houses, garbage cans, and manure piles. Avoid contamination of feed and foodstuffs. Repeat as necessary.

NOTICE

CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep container closed and away from heat, sparks, and open flame. Do not use in commercial food preparation areas or in edible products areas of food processing plants. Do not reuse empty container; bury or discard in a safe place. Destroy by perforating or crushing. Keep out of reach of children. This product is toxic to fish. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label.

CONTENTS: FIVE GALLONS

USDA Registration No. 2724-60

APPROVED FOR
DAIRY BARN



SUPER-M

50%

MALATHION INSECTICIDE SP


A multipurpose insect spray that controls:

Flies, Mosquitoes, Aphids, Red Spiders, Leafhoppers, Certain Species of Beetles, Caterpillars, and Certain Other Insects.

CAUTION: KEEP OUT OF REACH OF

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS: Malathion (0, 0-Dimethyl dithiophosphate of succinate) 50%; Aromatic petroleum hydrocarbons 40%;
INERT INGREDIENTS 10%





DEODORIZED
FORMULA

UPER-MAL

50%

MALATHION INSECTICIDE SPRAY

multipurpose insect spray that controls:

Flies, Mosquitoes, Aphids, Red Spiders, Thrips,
Leafhoppers, Certain Species of Beetles,
Caterpillars, and Certain Other Insects.

CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

INGREDIENTS: Malathion (0, O-Dimethyl dithiophosphate of diethyl mercapto-
) 50%; Aromatic petroleum hydrocarbons 40%;
INGREDIENTS 10%

DIRECTIONS FOR USE

FLOWERS, SHRUBS AND SHADE TREES: Use at the rate of 1 quart per 100 gallons of water (2 teaspoonfuls per gallon). Mix well and spray plants thoroughly. This spray will kill aphids, red spider mites, bagworms, leaf-hoppers, thrips, scales, mealybugs, white flies, leaf miners, leaf-bugs, adult Japanese beetles, lace bugs, plantbugs and tent caterpillars. Repeat treatment as necessary. Injury may occur to certain tender plants such as certain ferns and some species of crassula. Do not treat these.

HOUSEHOLD INSECTS: For control of Roaches, Ants, Spiders, Silverfish, Crickets, Clover Mites and Earwigs, use at the rate of 3 tablespoonfuls per quart (1 quart per 5 gallons) of water. Apply as a coarse spray or with a paint brush to window sills, baseboards, drainboards, under sinks and stoves and other areas frequented by insects.

FRUIT TREES: Use at the rate of 2 teaspoonfuls per gallon of water. Mix well and spray plants thoroughly. Repeat as necessary. This spray will kill the following:

ON PEACHES: Spider mites, plum curculio, Oriental fruit moth. Do not apply later than 7 days before harvest.

ON PEARS: Codling moth, pear psylla, plum curculio, leaf-roller. Do not apply later than 3 days before harvest. May cause injury to Bosc pear under certain conditions in the Northeast.

ON APPLES: Codling moth, plum curculio, spider mites, aphids, bud moth, Forbes scale, leaf roller. Do not apply later than 3 days before harvest. May cause injury to McIntosh varieties in summer sprays.

VEGETABLES: Use at the rate of one teaspoonful per gallon of water on the following vegetables. Mix well and spray plants thoroughly. **TOMATOES:** Aphids, and spider mites. **PEAS:** Aphids. **BEANS:** Leafhopper, Mexican bean beetle, spider mites. **PEPPERS and RUTABAGAS:** Aphids. **BROCCOLI:** Aphids, cabbage worm, cabbage looper. **EGGPLANT:** Aphids, spider mites. **CUCUMBERS, SQUASH and MELONS:** Aphids, spider mites. **ONIONS:** Onion thrips. **POTATOES:** Aphids, leaf hoppers. On these crops do not apply later than 3 days before harvest. Do not apply to cucurbits unless plants are dry.

Use at the rate of 2 teaspoonfuls per gallon of water on the following: **LETTUCE:** Aphids, leafhoppers. Do not apply later than 14 days before harvest. **SPINACH:** Aphids. **BEETS and BRUSSEL SPROUTS:** Aphids. Do not apply later than 14 days before harvest.