

DIRECTIONS FOR USE

DAIRY, BEEF CATTLE AND HORSES  
LIVESTOCK FLY CONTROL

**FACE FLIES**—For the control of Face Flies, Horn Flies, and Stable Flies, dilute 1 gallon of DDT (100% DDT) with 7 gallons (1 quart per gallon) of water and apply to all parts of the animal thoroughly. Repeat as necessary but not more often than once weekly.

**FACE FLIES**—For the control of Face Flies, dilute 1 gallon of DDT (100% DDT) with 7 gallons (1 quart per gallon) of water and apply to all parts of the animal thoroughly. Repeat as necessary but not more often than once weekly.

**HORN FLIES**—For the control of Horn Flies dilute 1 gallon of DDT (100% DDT) with 14 gallons (1 quart per gallon) of water, and apply by pouring 1 quart of the diluted emulsion along the backline of the animal. Repeat as necessary but not more often than once weekly.

**BACK RUBBER**—For the control of Horn Flies on Dairy and Beef cattle, dilute 1 gallon of DDT (100% DDT) with 18 gallons of fuel oil or other suitable carrier and charge backrubber device or sack or cloth as occasion requires. Retreat backrubber as needed. Do not spray animals with this oil solution.

**LICE AND TICKS**—For the control of Lice and Ticks on Cattle, Sheep, Goats, Horses and Dogs dilute 1 gallon of DDT (100% DDT) with 10 gallons of water and spray animal thoroughly. Make second application in 14 days if needed. Repeat as necessary but not more often than once weekly.

**BARN FLY CONTROL**—As an aid in the solution of House Flies, Face Flies, and Stable Flies in dairy barns, and in stables, hog barns and poultry houses, and other places where flies are a problem, mix 1 gallon of DDT (100% DDT) with 2 gallons (1 pint per gallon) of water. Spray mixture on surfaces where flies alight and feed. Spray once to remove flies, but not more than one gallon of spray per 1000 sq. feet. To improve control add 1 pound of kerosene to each gallon of spray solution.

**WARNING**—Hazardous if swallowed. Irritated on absorbed through skin. Irritates on skin, on clothes. Wash with soap and water after use. Avoid contamination of food, water, and food storage containers. Avoid contact of product with exposed skin, clothes and lead in operation. Wear clean synthetic rubber gloves and a cap or respirator. Do not use in areas controlled by the U.S. Department of Agriculture for protection against crop-damaging insects. Keep away from food and especially away from children and babies. Do not use on containers. Do not use on sheep, do not use on material of which a person wears an eye. Do not use on cattle.

**INTERNAL**—Internal—In case of poisoning on person or clothing, immediately remove clothing and flush skin or clothing with water. Call a physician for medical attention.

**INTERNAL**—Call a physician immediately. Symptoms—Nausea, vomiting, dizziness. Antidote—Activated charcoal. Do not induce vomiting.

NET CONTENTS: ONE GALLON

DAIRY, BEEF CATTLE AND HORSES

7-11-67

2724-101

**Warning:** May be fatal if swallowed, inhaled, or absorbed through skin. Do not breathe spray mist. Do not get in eyes or on skin. Wash hands, arms, and face with soap and water after handling and before eating or smoking. Rapidly absorbed through skin. Wash all contaminated clothing with soap and hot water before reuse. Wear natural rubber gloves. Avoid contact with eyes and mouth of animal. Do not apply to areas which will be contacted frequently by children and do not allow them in treated areas until surfaces are dry. Do not contaminate food, water, food containers, or cooking utensils.

**Antidote:** External — wash with soap and water; Internal — call a physician immediately. Large doses of Atropine are antidotal. Give emetic of mustard.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label instructions.

U.S.D.A. REG. NO. 2724-65

SOLD BY  
**VET-KEM LABORATORIES INC.**  
A THURON COMPANY  
12200 DENTON DRIVE, DALLAS TEXAS 75234

© 1970 Vet Kem Laboratories Inc

**vet-kem**

FOR USE BY OR UNDER THE SUPERVISION OF VETERINARIANS ONLY

**kem-dip**

Warning:

**vet-kem**

**kills resistant ticks & fleas**

**ACTIVE INGREDIENTS:**

Dioxathion 2,3-p-dioxanedithiol S,S-bis (O, O-diethyl phosphorodithioate) 3.50%; F  
1.50%; Diethyl diphenyl dichloroethane 9.50%; Related compounds 0.50%; 2,2-di  
phosphate 0.93%; Related compounds 0.07%; Pine oil 25.00%; Petroleum distill  
**INERT INGREDIENTS: 10.00%.**

**NET CONTENTS: 1 GALLON**

**WARNING: KEEP OUT OF REACH OF CHILDREN**  
SEE BACK PANEL FOR ADDITIONAL WARNINGS

antidote

**Directions for use**

**For Control of Ticks & Fleas on Dogs:** Dilute **1** gallon of KEM-DIP with **25 to 50** gallons of water (**2½ to 5** ounces per gallon) and dip, sponge or spray dogs thoroughly. Repeat as necessary, but not more often than once weekly.

**For Control of Ticks & Fleas in Doghouses, Lawns, & Outdoor Areas:** Dilute **1** gallon of KEM-DIP with **10** gallons of water (**12** ounces per gallon) and spray infested areas. Pay particular attention to places where dogs have been sleeping or resting for the past several months. Repeat treatment as necessary. Do not dip dogs with this solution.



PRODUCT LABEL FOR: KEM-DIP, KILLS RESISTANT TICKS & FLEAS

PL-07249-B

Apr. 13, 1971

2724-B

2724-B