

ZOECON RF-32 LIVESTOCK SPRAY & DIP

and backrubber refill

(contains dioxathion and dichlorvos)

Controls Hornflies, Lice & Ticks

NO WAITING PERIOD BEFORE SLAUGHTER

ACTIVE INGREDIENTS:

Dioxathion: [2,3-p-Dioxanedithiol S,S-Bis (0,0-Diethyl phosphorodithioate)]....	10.50%
Related Compounds.....	4.50%
2,2-Dichlorovinyl dimethyl phosphate.....	0.50%
Related compounds.....	0.04%
INERT INGREDIENTS:.....	84.46%


**KEEP OUT OF REACH OF CHILDREN
WARNING**

See back panel for additional precautionary statements

NET CONTENTS:

EPA Reg. No. 2724-80

EPA Est. 2724-TX-1



PRECAUTIONARY STATEMENTS

RM 15
File # 272-577

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - WARNING: May be fatal if swallowed, inhaled or absorbed through skin or eyes. Do not get in eyes or on skin. Do not breathe fumes or spray mist. Do not contaminate food or feedstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash contaminated clothing before reusing. Do not treat animals under 3 months of age.

STATEMENT OF PRACTICAL TREATMENT: If swallowed - Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE** vomiting. Vomiting may cause aspiration pneumonia. If inhaled - Remove victim to fresh air. Apply respiration if indicated. If on skin - Wash immediately with soap and water. If in eyes - Flush eyes with plenty of water. Call a physician immediately.

NOTE TO PHYSICIAN: If signs of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds and wildlife. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL HAZARDS: Do not use or store near heat or open flame.

DIRECTIONS FOR USE - It is a violation of Federal law to use this product in a manner inconsistent with its label.

DO NOT USE ON DAIRY ANIMALS OR IN DAIRY BARNS

Do not treat animals under three months of age.

On Cattle and Horses: To control Horn Flies, Cattle Lice, Ear Ticks, Lone Star Ticks, Gulf Coast Ticks, Winter Ticks and Cattle Fever Ticks, mix 1 quart of Zoecon RF-32 Livestock Spray & Dip with 25 gallons of water and stir well to form a creamy emulsion. Treat animals with a strong spray to penetrate to the skin, or dip thoroughly. Hand dip young animals. Approximately two quarts of spray are required for adult animals and proportionately less for younger animals. Be sure not to overdose young animals. For Ear Ticks and Gulf Coast Ticks, apply with low pressure to ears, head, neck and shoulder regions. For Tick control, repeat as necessary but not more often than once every two weeks. One spraying of Zoecon RF-32 Livestock Spray & Dip will control Horn Flies up to three weeks and a single treatment will often control Lice. Repeat as necessary, but not more often than every 2 weeks. Prior to dipping, water animals well to prevent drinking of vat fluid or runoff.

On Sheep and Goats: To control Sheep Lice, Goat Lice, Sheep Ticks (Keds), and Wool Maggots, use 1 quart of Zoecon RF-32 Livestock Spray & Dip with 25 gallons of water, mix well and spray or dip animals thoroughly. When dipping animals, be sure to wet them thoroughly. Always stir dipping vat well just before using. Repeat as necessary, but not more than every 2 weeks.

On Hogs: To control Hog Lice, use 1 quart of Zoecon RF-32 Livestock Spray & Dip to 25 gallons of water, mix well, and spray animals with a strong spray to thoroughly wet the skin. Repeat as necessary, but not more often than every 2 weeks. Do not treat sows within 2 weeks of farrowing or while nursing.

In Backrubbers: As an aid in the control of Horn Flies and Lice on beef cattle, dilute 1 quart of Zoecon RF-32 Livestock Spray & Dip with 9 quarts of fuel oil or other suitable carrier and charge backrubber device or sock sack or cloth as occasion requires. Retreat backrubber as needed. Do not spray animals with this solution. Do not "pour on" oil solutions.

ZOECON RF-32 LIVESTOCK SPRAY & DIP DILUTION TABLE

AMOUNT OF ZOECON RF-32	GALLONS OF WATER		VOLUME OF OIL	
	SPRAY OR DIP	BACK-RUBBER	SPRAY OR DIP	BACK-RUBBER
1 Quart	25 Gallons	9 Quarts	25 Gallons	9 Quarts
1 Gallon	100 Gallons	36 Gallons	100 Gallons	36 Gallons

STORAGE & DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Plastic container:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by procedures approved by state or local authorities. **Glass container:** Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state or local procedures.

ZOECON INDUSTRIES
A Division of Zeecon Corporation
12200 Denton Drive, Dallas, Texas 75234

Made in USA

© 1981 Zeecon
184-B-0948-A-0088A