

ACCEPTED
APR 4 - 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT GIDE ACT
FOR ECONOMIC POISON REGIST
ED UNDER NO. 2724-78 SUBJECT
TO ATTACHED COMMENTS.

STARBAR
VAPONA[®] MIST
FOR FOGGING OR SPRAYING

For use in Foggers, Mist Sprayers and Hand Sprayers. Ready to use solution for the control of House Flies, Mosquitoes, Gnats, Wasps, ~~Fleas, Ticks, Face Flies, Horn Flies, Stable Flies in Outdoor areas,~~ Patios, Restaurants, Warehouses, Food Processing Plants, Horse Barns, Dairy Barns, Shelter Sheds, Milk Sheads.

ACTIVE INGREDIENTS:

| | |
|--------------------------------------|----------------|
| 2,2-dichlorovinyl dimethyl phosphate | 0.93% |
| related compounds | 0.07% |
| Mineral oil | 99.00% |
| | <u>100.00%</u> |

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

Net Contents: One fluid quart

DIRECTIONS FOR USE

VAPONA MIST MAY BE USED TO TREAT: OUTDOOR LIVING AREAS, PICNIC GROUNDS, BACKYARD AREAS, PATIOS, LOADING DOCKS, OUTDOOR LATRINES, PARKING AREAS, REFUSE AREAS, AROUND SERVICE STATIONS, OPEN-AIR DRIVE-INS, ICE CREAM STANDS, GARBAGE COLLECTION AND DISPOSAL AREAS, RESTAURANTS, THEATERS, FOOD PROCESSING PLANTS, INDUSTRIAL PLANTS AND WAREHOUSES.

CATTLE (Beef and Dairy) to control Hornflies, Houseflies, Stable Flies, Face Flies, apply 1 or 2 fluid ounces of Vapona Mist per animal as a mist spray daily with hand or automatic sprayer. Apply spray so as to wet the hair. DO NOT apply in excess of 2 fluid ounces per animal per day.

ANIMAL BUILDINGS (Horse Barns, Dairy Barns, Shelter Sheds) Mild Sheds. Outdoor areas. To control Flies, Gants, Mosquitoes, Roaches, Fleas, Ants, Sowbugs, Wasps, Flying Moths, and other small flying insects. Cocroaches, Ants, Spiders and Silverfish.

~~Apply Vapona Mist as a coarse wet spray at the rate of 1 pt. of 1.0% solution per~~
1,000 sq. ft. for milk sheds, stables, livestock barns, loafing sheads, pig pens, and outdoor areas, treat around doorways, feed storage rooms, alleyways and window sills. DO NOT contaminate milk or milking utensils. For control of Fly Maggots in Maggot breeding areas apply at the rate of 1 to 2 pts. per 100 sq. ft.

IN POULTRY HOUSES: Apply to manure, window sills outside of penned enclosures and on floor of feed storage rooms.

IN DOG KENNELS: Apply to outside dog runways, window sills, and ledges. For control of fleas and brown dog ticks increase dosage to 1 qt. of Vapona Mist per 1,000 sq. ft. Treat infested areas, baseboards, cracks, walls, doors. DO NOT allow children in treated area until dry.

For control of fly maggots in maggot breeding areas, apply at the rate of 1 to 2 pts. per 100 sq. ft.

FOR FOGGING: Use 1 pint of Vapona Mist per 8,000 cu. ft. or 1 gallon per every 50,000 cu. ft. indoors. When fogging indoors, eliminate all sources of ignition (pilot-lights, fires, open flames) and disconnect electric power. Apply dense fog when wind is calm 1 to 5 seconds per 1000 su. ft. Before applying, reduce air movement as much as possible by closing doors, windows and other openings. DO NOT contaminate water, feed or foodstuffs, milk or milking utensils. DO NOT use this solution for fogging purposes in areas where animals have received direct application within 8 hours.

Occupants should vacate premises before indoor treatments and should not reoccupy the premises until it has been ventilated. After application open several windows and doors to create air currents to ensure reduction of Insecticide vapors. Food should be removed and food handling equipment covered with wrapping paper during fogging or mist spraying indoors, use mask or respirator of a type passed by the U. S. Department of Agriculture for DDVP protection. DO NOT FOG INTO OR NEAR AN OPEN FLAME.

2724-P

ACCEPTED
APR 4 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT

ACCEPTED WITH COMMENTS

DIRECTIONS FOR USE (cont'd)

CAUTION: May be fatal if swallowed, inhaled or absorbed through the skin or eyes. Wash thoroughly with soap and water after handling and before eating or smoking. DO NOT contaminate feed, water or foodstuffs. Keep out of the reach of children. If swallowed, induce vomiting immediately and get medical attention. Do not get in eyes, or skin or on clothing. Rapidly absorbed through skin or eyes. Do not breath vapor mist. In case of contact, immediately remove contaminated clothing and flush skin and eyes with plenty of water: for eyes get medical attention. Wear clean rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for protection against DDVP. Wash contaminated clothing with soap and hot water before reuse. Do not reuse shoes or boots until free of all contamination. ATROPINE is antidotal. Do not contaminate any body of water by the cleaning of equipment or disposal of wastes and containers. Toxic to aquatic organisms and fish.

U. S. D. A. Reg. No. 2724-98
STARBAR DIV.
THURON INDUSTRIES, INC.
12200 Denton Drive
Dallas, Texas 75234

