

— 10 —

Volume 106

Kidney Disease, Malaria, & other Child Diseases, and  
their up to 3 month's

## Decomposition Induction Method

USE IN HOME/MONITOR/MICROPHONE UNIT

**CAUTION:** Keep out of reach of children.  
See back page for additional information.

## CONTINUATION OF THE HISTORY OF THE CHURCH IN THE UNITED STATES.

## Minimizing the cost of energy use

100%  
D.D.T.

VET-KEN™

VAPOR BAR

DO NOT OPEN UNTIL FLYING INSECTS ARE PRESENT


To control flies, mosquitoes, gnats and other small flying insects and silverfish.

Use in homes, motels, dairy barns, milk rooms and animal shelters.

DIRECTIONS FOR USE

Homes - Motels

To control flies, mosquitoes, gnats, other small flying insects and the reduction of silverfish, place one Vet-Ken Vapor Bar per average size room (10 x 10 x 10 feet) or per 100 square feet of floor area. Normally this is effective for three months. Replace strips if effectiveness diminished. For control of flies in garbage cans attach one Vet-Ken Vapor Bar inside of an average size garbage can (20 gallon capacity). One unit will be effective for the entire fly season. Keep lid on garbage can.



Agricultural Areas

To control flies, mosquitoes, gnats and other small flying insects in Horse barns, Dairy Barns, Milk Barns, Milk Sheds, Stables, Milk rooms, Pie Plants, Dog Kennels, poultry houses and other farm buildings, suspend one Vet-Ken Vapor Bar (1 1/4" x 3 7/8" x 1 1/4") for each 100 square feet of floor space. If more than one bar is required, distribute them so that no area to be treated by either land or water is left untreated. The other uses mentioned in the directions are also effective for outdoor areas. Do not open until flying insects are present.

Catch Fly Trap

For the control of flies and gnats place two suspended Vet-Ken Vapor Bars in a catch fly trap and open the trap. Place the trap near the flies. Place the trap about 10 feet from the house or building where the flies are most numerous.

NOTES

Caution: This product contains DDT which is a potent poison. Do not eat or drink while handling this product. Wash hands after handling. Avoid contact with skin and eyes. Do not smoke while handling this product. Do not store in direct sunlight. Store in a cool dry place.

placed in hospital or clinic rooms, with an patient room, parlor, nursery, operating or emergency areas. Do not use in any room where infants, the sick, or aged are or will be present for any extended period of confinement. Do not use in kitchens, laundry, etc., or areas where food is prepared or served.

**MERRILL:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the Label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to Label instructions.

The of the packaged product is licensed under Shell's U.S. Patent 2,956,070 and 3,116,201 for use as an insecticide generator only. Not to be taken internally by humans or animals.

ACTIVE INGREDIENTS

## *Z,Z-dichlorovinyl diacetyl phosphide* compounds.

116

## Table I. Summary

1. *Leucosia* *leucostoma* *leucostoma* *leucostoma* *leucostoma* *leucostoma*