

STARBAR



SIMAX[®]

DAIRY AND BEEF CATTLE DUST
CONTAINS 3% CIODRIN* INSECTICIDE

CONTROLS HORNFLIES AND LICE

APPROVED FOR DIRECT APPLICATION TO LACTATING
DAIRY CATTLE. WILL NOT CONTAMINATE MILK WHEN USED AS DIRECTED.

ACTIVE INGREDIENTS:

dimethyl phosphate of alpha-methylbenzyl
3-hydroxy-cis-crotonate 3.0%

INERT INGREDIENTS: 97.0%

100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS



Sold by
STARBAR, INC.
A Thuron Company
12200 Denton Drive
Dallas, Texas
75234

Net Contents:
Four Pounds Avdp.



SIMAX[®] DAIRY AND BEEF CATTLE DUST

...may show hypersensitivity to organic phosphates.

FOR DIRECT APPLICATION TO DAIRY AND BEEF CATTLE TO CONTROL HORNFLIES AND LICE: For the control of hornflies on beef and lactating dairy animals, apply 1 - 2 heaping tablespoonsful to the backline of the animal, covering in addition the poll, neck, and upper portions of the sides. Repeat above application as necessary to maintain control, but not more often than once every 14 days. Wash hands thoroughly after use.

For the control of lice on dairy and beef cattle, treat animals thoroughly. If required, repeat this thorough application in 3 - 4 weeks.

DUST BAG: For control of hornflies, and as an aid in the control of lice, on dairy cattle and beef cattle, SIMAX DUST may be applied daily when application is made with Dust Bags. To prepare dust bags for animal use, place 4 - 8 pounds of SIMAX DUST in cotton cloth or double burlap bags and suspend bags near mineral or salt blocks, in alleyways leading to or from animal building, or dairy barns. Dust bags may also be suspended in livestock housing pens, feed lots, and loafing sheds where possible. Protect bags from weather by placing them under shelter.

CAUTION: AVOID PROLONGED OR REPEATED inhalation of dust and contact with skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Keep away from open flame. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, or foodstuffs. Do not contaminate any body of water, by direct application, cleaning of equipment, or disposal of wastes or containers. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label instructions.

U.S.D.A. Reg. No. 2724-146

*Ciadrin Reg. Trademark Shell Chemical Co.

Sold by
STARBAR, INC.
A Thuron Company
12200 Denton Drive
Dallas, Texas 75234

SIMAX[®]
DAIRY AND BEEF
CATTLE DUST
NET WT. 4 LBS.



vet-kem[®]

AVAILABLE THROUGH VETERINARIANS

a
beef
cattle
insecticide
for
control
of
cubs,
lice,
and
botflies

kemolate

emulsifiable liquid

Directions for use: ...
Dose: ...
Precautions: ...
Warnings: ...

WARNING: KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. ...
Do not use in areas where children play. ...
Avoid contact with eyes, nose, and mouth. ...