

flea collar for dogs

Do not open until ready for use.

CAUTION: Do not allow children to play
with this collar.

ACTIVE INGREDIENTS:

2, 2-dichlorovinyl dimethyl phosphate 8.37%

Related Compounds 0.63%

INERT INGREDIENTS: 91.00%

For full cautions and directions for use,
see accompanying carton label.


P454-B

LOT NO H3G90 CR

MADE BY VET-KEM LABORATORIES INC. A THURON COMPANY
12200 DENTON DRIVE DALLAS TEXAS 75234



vet-kem



Directions for use

Buckle Flea Collar around dog's neck. Collar must be worn loosely. Any collar when buckled too tightly may cause skin irritation. Attach collar so that two or three fingers may be inserted easily between collar and dog's neck. Cut off any excess length of collar and dispose of it. The Flea Collar starts killing fleas as soon as it is placed around the dog's neck. The fleas on the dog will be killed and new ones which may temporarily appear on the dog will also be killed during the 3 months (13 weeks) collar is worn. TICKS are tough and are killed slowly. They will usually be dead in 1-5 days. Collar will continue to kill ticks for 8 weeks. Dead ticks may occasionally remain in dog's coat but are easily detached. Ticks inside ears are hardest to kill — for best results remove these ticks when collar is put on. Use in addition to regular collar. When bathing dog remove collar until dog is dry. Replace collar when effectiveness diminishes.

ACTIVE INGREDIENTS

2,2-dichlorovinyl dimethyl phosphate *8.37%
Related Compounds *0.63%

INERT INGREDIENTS 91.0%

Caution: Do not open inner envelope until ready to use. Do not allow children to play with this collar. Collar is intended for use only as an insecticide generator (Shell's U.S. Patents 2,956,073 and 3,116,201) and is not to be taken internally by man or animals. Do not use on cats, whippets and greyhounds. Some dogs may be sensitive to this collar. Animals should not be allowed to chew on the collar. When collar is first worn, observe neck area for irritation every few days. Remove collar at the first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs. Do not use any other pesticide on dog while collar is worn.

*Equivalent to 9.0% DDVP insecticide

kemic flea collar

flea collar



vet-kem



kemic


for dogs – black

CONTENTS: 1 COLLAR – 2/3 OZ NET WT

CAUTION: DO NOT ALLOW CHILDREN TO PLAY WITH THIS COLLAR
SEE CAUTIONS ON BACK

© 1969 Vet-Kem Laboratories, Inc.

16027-B



27-29-111

7

flea collar for dogs

Do not open until ready for use

CAUTION: Do not allow children to play with this collar.

ACTIVE INGREDIENTS

... ..

INERT INGREDIENTS

... ..

For full instructions, see accompanying carton label.

LOT NO H3390 CR