


SMAX

A NEW CHEMICAL FOR EFFECTIVE RODENT CONTROL
CONTAINS PROLIN®



INSTRUCTIONS FOR USE

After opening SMAX BAIT STATION place it in a protected spot where the rats and mice ordinarily feed and water, or where they travel or live. SMAX Bait Station should be placed along walls and in corners rather than in the middle of the room.

If this container is not used as a Bait Station, pour SMAX into shallow container. **CAUTION**—Place baits in locations not accessible to children, pets, and domestic animals or in tamper proof bait boxes.

FOR RATS: Use one or two bait stations at every point of large activity. Look for such tell-tale signs as gnawings, tracks or trails, burrows, droppings or damage.

FOR MICE: Use a larger number of smaller bait placements——about 1 tablespoon each at 8-10 foot intervals.

KEEP FEEDING STATIONS WELL SUPPLIED

It is important to inspect and replenish each station daily for the first week and regularly thereafter until bait is no longer being taken. Never allow a feeding station to remain empty for over 48 hours. If some bait stations are untouched, move to a different location. If baits become sour or moldy (as might be the case during extended periods of hot humid weather), replace with fresh bait as needed.

Where re-infestation from dumps and fields may occur, permanent SMAX Bait Stations should be set out and replenished as needed.

CAUTION

If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions may be indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

A PRODUCT OF STARBAR, INC.
DIV. THURON INDUSTRIES, INC.
DALLAS, TEXAS
U.S. D. A. Reg. No. 2724-N

ACCEPTED
MAY 24 1967
UNDER FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC CONTROL REGISTERED
ED UNDER NO. 2724-1 SUBJECT
TO ATTACHED CONDITIONS.

SEARBAF
MAX
RAT AND MOUSE BAIT
BAIT STATION

ACTIVE INGREDIENTS:

Warfarin [3(a-acetonylbenzyl)-4-hydroxycoumarin]	0.025%
N ¹ -(2-quinoxaliny) sulfamylamide (Sulfaquinoraline)	0.025%

INERT INGREDIENTS:

	99.950%
Total	100.000%

CAUTION: KEEP OUT OF REACH OF CHILDREN
See Back panel for additional cautions.

NET WEIGHT:
1 POUND AVDZ.

