


16092



STAGBAR

HOG MANGE AND LICE SPRAY

Combining Malathion & Toxaphene for Quick Kill & Long Residual

ACTIVE INGREDIENTS: Toxaphene (Technical Chlorinated Camphene) 67.5% Malathion (0, 0-Dimethyl Diminophosphate of Methyl S-methylpropyl ester) 32.5% Petroleum Distillate 0.7%

INERT INGREDIENTS: 32.8%

NOTE: This is a contact spray. Apply to all surfaces to be treated. Residual activity is long.

Net Weight 1.00

Net Weight 1.00

For further information, contact the manufacturer.

2( of 2)

DIRECTIONS FOR USE

SPRAY - Effective against sarcoptic mites and hog lice on swine, dilute 1 ounce of SPRAY in 100 gallons of water, mix well and spray thoroughly. One thorough treatment is sufficient.

DIP - Effective against sarcoptic mites and hog lice on swine, dilute 1 ounce of HOG MANE or SPRAY to 100 gallons of water, mix well and dip animals thoroughly. One thorough treatment is sufficient.

**WARNING:** Avoid treating during cold, stormy weather; avoid treating overheated or sick animals; water animals before dipping. Do not treat animals under three months of age. Individual dipping of young animals is recommended to avoid swallowing of dipping solution.

May be fatal if swallowed. Do not apply in dairy barns or poultry houses. Avoid contamination of feed and foodstuffs. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this spray or washings from the spray tank. Do not use on animals within 28 days of slaughter. Do not treat cows within 2 weeks of milking nor for at least 3 weeks thereafter.

**REMARKS:** May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or clothing.

DILUTION TABLE

<u>VOLUME OF HOG MANE OR SPRAY</u>	<u>VOLUME OF WATER</u>
1 PINT	100 GAL.
1 QUART	25 GAL.
1 GALLON	10 GAL.

ACCEPTED WITH COMMENTS

124-157  
LACY