

1 (of 2)

STARBAR
®

LINTOX-M DUST

For Beef Cattle, Sheep, Goats and Hogs

Active Ingredients:

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate)	4%
Toxaphene (Chlorinated camphene contain- ing 67 to 69 percent chlorine)	5%

Inert Ingredients: 91%

CAUTION: Keep out of reach of children.
See back panel for additional cautions.

SOLD BY

STARBAR, INC.
A THURON COMPANY
12200 Denton Drive - Dallas, Texas 75234

NET CONTENTS: 25 POUNDS

ACCEPTED
JUN 5 1969
UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 2724-164 SUBJECT
TO ATTACHED COMMENTS.

2724-164

ACCEPTED
Jan 5 1969

2 (of 2)

Directions for Use

BEEF CATTLE

For the control of Hornflies, Ticks and Lice on beef cattle, treat animals thoroughly and rub dust lightly into hair. Repeat treatment as needed. Do not apply within 30 days of slaughter.

SWINE

For control of Hog Lice, apply dust to hogs, pens, and litter. One treatment may be sufficient. Treat animals thoroughly. Repeat in 10 days or thereafter as needed. Do not apply within 30 days of slaughter.

SHEEP & GOATS

For control of Lice and Keds on sheep and goats, treat animal thoroughly preferably following shearing. Repeat as needed. Do not apply within 30 days of slaughter. Do not apply to milk goats.

CAUTION

Harmful if swallowed. May be absorbed through the skin. Avoid prolonged breathing of dust. Avoid contact with eyes, skin and clothing. Avoid prolonged or repeated contact with skin, wash thoroughly after using. Do not use on lactating dairy cattle, in dairy barns or on calves under three (3) months of age. Do not contaminate feed or foodstuffs. Do not apply within 30 days of slaughter. Do not apply to milk goats. Avoid storage near feed and food products.

This product is considered toxic to fish and wildlife. Do not contaminate streams, lakes, or ponds.