



4 oz. carton label

AMENDED
MAR 15 1978
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 2724-169

Do not give anything by mouth to an unconscious person. Call a physician or Poison Control Center immediately.

Do not get in eyes, on skin or clothing. Do not breathe spray mist. Use only in well-ventilated areas. Wear rubber gloves, goggles and protective clothing. In case of skin contact, wash immediately with soap and water, for eyes, flush with water. Wash all contaminated clothing with soap and hot water before re-use. Do not store near heat or open flame. Do not contaminate food or feed.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Container disposal: Do not use empty container. Triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, bury in a safe place or wrap container and put in trash collection.

Note to physician and veterinarian: This is an organophosphorus insecticide. Atropine is antidotal. Usual symptoms of organophosphorus poisoning in man include headache, blurred vision, weakness, nausea, discomfort in the chest, vomiting, abdominal cramps, diarrhea, salivation, sweating, and pin-point pupils. Usual symptoms of organophosphorus poisoning in animals include salivation and labored breathing.



vet-kem®

AVAILABLE THROUGH VETERINARIANS

kemolate™
emulsifiable liquid

An insecticide for the control of sarcoptic mange on dogs, and for fleas and ticks on dogs and cats

WARNING: KEEP OUT OF REACH OF CHILDREN
 READ ALL DIRECTIONS AND PRECAUTIONS BEFORE USING. See side panel for note to physician and veterinarian.

CONTENTS: 4 FL. OZ.

SAVE THIS CARTON FOR FULL DIRECTIONS AND PRECAUTIONS

Gives control of sarcoptic mange on dogs with one dip treatment

Gives 16 day residual control of fleas and Brown Dog ticks on dogs

Gives 9 day residual control of fleas on cats

Gives 16 day residual control of ticks on cats

ACTIVE INGREDIENTS:

N-(Mercapto-methyl) phthalimide 11.60%, Aromatic petroleum solvent 72.90%

INERT INGREDIENTS 15.50%

Protect from temperatures below 20 F

EPA Reg No 2724-169

EPA Est 2724-TX-1

US Pat No 2,767,194

A Division of Zoecon Corp
c 1978 Zoecon
Made in USA

ZOECON INDUSTRIES
12200 DENTON DRIVE
DALLAS TEXAS 75234

Directions: Sarcoptic mange on dogs is characterized by loss of hair, thickened skin and intense itching, caused by the mite *Sarcoptes scabiei var canis*. Mix 1 oz (2 tbs.) Kemolate with 1 gal water. Dip dog until skin is wet and allow to shake dry. Do not rinse. If no improvement is seen within 14 days, consult your veterinarian. If mange mites reinfest your dog, retreatment may be necessary.

Fleas and ticks on dogs: Mix 1 oz (2 tbs.) Kemolate with 1 gal water. Dip or sponge on solution until skin is wet. For maximum residual control, allow to dry on animal.

Occasionally, cats get ticks.

Fleas and ticks on cats: Mix 1/2 oz (1 tbs.) Kemolate with 1 gal water. Dip or sponge on solution until skin is wet. For maximum residual control, allow to dry on animal.

Retreat as necessary but not more often than every 7 days. Do not treat dogs or cats under 8 weeks of age. The Kemolate solution may be stored or used for 30 days.

Observe the following use precautions.

Warning: Kemolate is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Do not treat sick or debilitated animals.

May be harmful if swallowed, inhaled or absorbed through skin. If swallowed, do not induce vomiting. Immediately give large quantities of water or milk. If vomiting does occur, give fluids again. Never

