

Directions for use

ACCEPTED

 AUG 1 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 224123 SUBJECT
 TO ATTACHED COMMENTS.

Buckle Flea Collar around dog's neck. Collar must be worn loosely. Any collar, when buckled too tightly, may cause skin irritation. Attach collar so that two or three fingers may be inserted easily between collar and dog's neck. Cut off any excess length of collar and dispose of it. The Flea Collar starts killing fleas when it is placed around the dog's neck. The fleas on the dog will be killed and new ones which may temporarily appear on the dog will also be killed during the 2 months (8 weeks) collar is worn. Use in addition to regular collar. When bathing dog remove collar until dog is dry. Replace collar when effectiveness diminishes.

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate) 9.00%

INERT INGREDIENTS: 91.00%

Caution: Do not open inner envelope until ready to use. Do not allow children to play with this collar. Do not use on cats. Some dogs may be sensitive to this collar. Animals should not be allowed to chew on the collar. When collar is first worn, observe neck area for irritation every few days. Remove collar at the first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs. Do not use any other pesticide on dog while collar is worn.

E.P.A. REG. NO 2724-173-AA-


*Reg. Trademark of Union Carbide Corp. for the active ingredient carbaryl

Thuron Industries, Inc 12200 DENTON DRIVE, DALLAS, TEXAS 75234
 MADE IN U.S.A.
 © 1971 Thuron Industries, Inc

 **Kemic Dust Flea Collar**



vet-kem®



Kemic Dust Flea Collar

Protect Flea Collar for Dogs

dust flea collar



Kemic Dust Flea Collar



DIRECTIONS FOR USE:

Buckle collar around dog's neck. This collar, like all collars, should be worn loosely to prevent irritation. When collar is first placed on dog, adult ticks will be killed in a few days and will fall off or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages, ticks are smaller than a match head and are difficult to see. The Starbar Tick Collar will also control these immature stages. If ticks are a problem, this collar should be worn constantly. **Replace when effectiveness diminishes.**

Fleas — This collar is also highly effective in preventing and killing fleas. As long as your dog is wearing a Starbar Tick Collar, do not use a separate flea collar. **Replace when effectiveness diminishes.**

ACTIVE INGREDIENT:

o-Isopropoxyphenyl methylcarbamate*9.4%

INERT INGREDIENTS:90.6%

CAUTION: Do not open protective pouch until ready to use. Do not allow children to handle this collar. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction. * In case of contact, flush eyes with water. Wash hands thoroughly with soap and water after handling collar. The dust released by this collar is a cholinesterase inhibitor.

Note to Physician: Atropine and *homatropine are antidotal.

When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Do not use on sick or convalescing dogs. Do not use other pesticides on dog while collar is worn. Remove collar when bathing dog.

EPA Reg. No. 2724-253.

Manufactured by: THURON INDUSTRIES, INC.
12200 Denton Drive — Dallas, Texas 75234

*U S Patent No 3,111,539

EPA Est. 2724-TX-1

Patent Pending



Tick Killing Collar for Dogs

Kills and prevents ticks Also kills fleas

For small and medium size dogs
with up to 21" necks

CAUTION: Do not allow children to play with this collar —
SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

CONTENTS: 1 Collar; Net Wt. 1 oz.

2724-253

DIRECTIONS FOR USE

Buckle collar around dog's neck. Collar should be worn loosely to prevent irritation. Attach collar so two or three fingers may be inserted easily between collar and dog's neck. Cut off any excess length and dispose of it. The collar starts killing fleas as soon as it is placed around dog's neck. Fleas on the dog will be killed and new ones which may temporarily appear on the dog will also be killed while collar is worn. Use in addition to regular collar. When bathing dog remove collar until dog is dry. This product has been shown to provide effective flea control for up to 21 weeks. **Replace collar when effectiveness diminishes.**

Ticks are tough and are killed slowly. When collar is first placed on dog, adult ticks will be killed over entire body in a few days and will fall off or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages, ticks are smaller than a match head and are difficult to see. The collar will also control these immature stages. Collar will have to be replaced more frequently for ticks than for fleas. This product has been shown to provide effective tick control up to 4 months. If ticks are a problem, this collar should be worn continuously. **Replace when effectiveness diminishes.**

ACTIVE INGREDIENT:

o-Isopropoxyphenyl methylcarbamate* 9.4%

INERT INGREDIENTS: 90.6%

CAUTION: Do not open protective pouch until ready to use. Do not allow children to handle this collar. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction.* In case of contact, flush eyes with water. Wash hands thoroughly with soap and water after handling collar. The dust released by this collar is a cholinesterase inhibitor.

Note to Physician: Atropine and homatropine are antidotal.

When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Do not use on sick or convalescing dogs. Do not use other pesticides on dogs while collar is worn. Remove collar when bathing dog.

EPA Reg. No. 2724-254


THURON INDUSTRIES, INC.

12200 Denton Drive, Dallas, Texas 75234

U.S. Patent No. 3,111,539

EPA Est. 2724-TX-1

Patent Pending



Dog Collar for Flea Control

Kills fleas up to 5 months
Kills ticks up to 4 months

CONTENTS: 1 Collar; Net Wt. 1 oz.

CAUTION: Do not allow children to play with this collar —
See cautions on back.