

1-11-72

KEMLINE FLEA FOAM FOR CATS

Controls Fleas on Cats & Kittens

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl Methylcarbamate).....	0.500%
Pyrethrins.....	0.050%
Technical piperonyl butoxide*.....	0.100%
N-octyl bicycloheptene dicarboximide.....	0.156%
INERT INGREDIENTS:.....	99.184%
Total.....	100.000%

*Consists of 0.08% (butylcarbityl) (6-propylpiperonyl) ether and 0.02% related compounds.

USE ABOVE 60°F.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Cautions

NET CONTENTS: 5 oz. MADE.

Shake well before using

Directions for use: Shake well and press button to release foam. Hold nozzle so it touches the hair and apply a strip of foam down the back, extending from the neck area to approximately one inch from the base of the tail. The foam will penetrate the hair, breaking down gradually to produce an effective back line coverage. This treatment will control fleas on cats and kittens. Repeat treatment if necessary, but not more than once weekly. Do not allow foam to get in eyes or on scrotum.

Caution: Do not contaminate feed, water, or foodstuffs. Keep out of reach of children. Wash hands after use. Contents under pressure. Do not store near sources of heat. Never puncture container or throw into fires or incinerators. Harmful if swallowed. Avoid contact with skin. Avoid breathing of vapors.

Do not treat kittens less than four weeks old.

Warranty: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label instructions.

Zoecon Corporation
A Sandoz Company
12200 Denton Drive, Dallas, Texas 75234

EPA Reg. No. 2724-177
0190-B:0157E

© 1970 Thuron Industries, Inc.
Made in USA

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.