FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO 2724: 225 SUBJECT
TO ATTACHED COMMENTS:



DIRECTIONS FOR USE

GOLDEN SUGAR BAIT is a fly bait formula which should be scattered over fly feeding surfaces to provide initial knockdown and reduction of the housefly population.

Use GOLDEN SUGAR BAIT in and around feed lots, poultry houses, livestock barns, dairy barns, refuse piles, kennels and other areas where Houseflies are a problem. Scatter bait at the rate of approximately 1/4 lb. per 500 sq. ft. of fly feeding area. Repeat application as necessary. Do not treat portions of buildings where poultry feed or where animals will lick.

CAUTION: Avoid contamination of feed and foodstuffs. May be harmful if swallowed. Store out of reach of children, poultry and domestic animals. Do not use inside homes where milk is stored or processed, or during milking time when milk or utensils are exposed. Wash thoroughly after handling. Wash all contaminated clothing with soap and hot water before reuse.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label instructions.

Thuron Industries, Inc. • 12200 Denton Drive • Dallas, Texas 75234

Made in U.S.A.

© 1973 Thuron Industries, Inc.

Hornoy. Gsta Gol Suga

ACTIVE INGREDIENT: Dimethyl (2 hydroxyethyl) phosphonate INERT INGREDIENTS:

CAUTION: Keep Ou See back panel for

E.P.A. Reg.

NET WEIGHT:

OBSERVE LAGE DIRECTIONS

FOR USE

t formula which should be scatprovide initial knockdown and

piles, kennels and other areas after bait at the rate of approxeding area. Repeat application of buildings where poultry feed word | Cennels

reach of children, poultry and e homes where milk is stored or when milk or utensils are exdling. Wash all contaminated ore reuse.

oy it by perforating or crushing. from water supplies.

and other wildlife. Keep out of ly where runoff is likely to occur. Instanton drift from areas treated. Ing of equipment, or disposal of pecified on this label.

concerning the use of this product other risk of use and handling of this material instructions.

nton Drive • Dallas, Texas 75234

© 1973 Thuron Industries, Inc.

homogenized Starbar

Golden Sugar Bait



Contains Trichlorfon

ACTIVE INGREDIENT: Dimethyl (2, 2, 2-trichloro-1-hydroxyethyl) phosphonate INERT INGREDIENTS:

0.40% 99.60%

CAUTION: Keep Out of Reach of Children
See back panel for additional cautions

E.P.A. Reg. No. 2724-225

NET WEIGHT: 1 POUND AVDP.