

Long Lasting
Long Lasting Protection

Kills Fleas Fast
Fast Acting for Fleas
Quick Killing for Fleas
Quick Acting for Fleas
Fast Action for Fleas

DIRECTIONS FOR USE

Buckle collar around dog's neck. Collar should be worn loosely to prevent irritation. Attach collar so two or three fingers may be inserted easily between collar and dog's neck. Cut off any excess length and dispose of it. The collar starts killing fleas as soon as it is placed around dog's neck. Fleas on the dog will be killed and new ones which may temporarily appear on the dog will also be killed while collar is worn. Use in addition to regular collar. This product has been shown to provide effective flea control for up to 21 weeks. **Replace collar when effectiveness diminishes.**

Ticks are tough and are killed slowly. When the collar is first placed on dog, adult ticks will be killed over entire body in a few days and will fall off or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages, ticks are smaller than a match head and are difficult to see. The collar will also control these immature stages. This product has been shown to provide effective tick control up to 5 months. If ticks are a problem, this collar should be worn continuously. Kills ticks that may carry and transmit Rocky Mountain spotted fever and tularemia. Temporary wetting will not reduce effectiveness. If your dog goes swimming or is out in the rain, it is not necessary to remove the collar. The continuous action collar will rapidly replace any flea and tick killing material removed. **Replace when effectiveness diminishes.**

ACTIVE INGREDIENT

o-Isopropoxyphenyl methylcarbamate 9.4%
INERT INGREDIENTS 90.6%

CAUTION: Do not open protective pouch until ready to use. Do not allow children to handle this collar. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction. In case of contact, flush eyes with water. Wash hands thoroughly with soap and water after handling collar. The dust released by this collar is a cholinesterase inhibitor.

Note to Physician: Atropine and homatropine are antidotes.

When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Do not use on sick or convalescing dogs. Do not use other pesticides on dogs while collar is worn.

EPA Reg No 2724-254
Zocon Industries
12200 Denton Drive, Dallas, Texas 75244
EPA Est 2724-TX 1

1280-B

ACCEPTED
MAR 24 1981
Under the Federal Insecticide, Fungicide and Rodenticide Act amended for the pesticide registration under FFA Reg No.

Dog Collar for Flea Control

Kills fleas up to 5 months
Kills ticks up to 5 months

From Head to Tail

Rain or Shine

CONTENTS: 1 Collar; Net Wt. 1 oz.

CAUTION: Do not allow children to play with this collar — See cautions on back

BEST DOCUMENT AVAILABLE