

THURON

TICK KILL COLLAR

For repackaging of an insecticide only – Not for resale

See attached sample label

Confidential data to support the label are on file with the Environmental Protection Agency in file EPA Reg. No. 2724-252.

ACTIVE INGREDIENT: <i>o</i> -Isopropoxyphenyl methylcarbamate*	9.4%
INERT INGREDIENTS:	90.6%

EPA REG. NO. 2724-255

CAUTION: DO NOT ALLOW CHILDREN TO HANDLE COLLAR

Do not open protective pouch until ready to use. Do not allow children to handle this collar. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction.* In case of contact, flush eyes with water. Wash hands thoroughly with soap and water after handling collar. The dust released by this collar is a cholinesterase inhibitor.

Note to Physician: Atropine and *homatropine are antidotal.

Do not reuse container. Destroy when empty.

NOTE: Additional data may be required to support registration of your product, if your labeling text or graphics are altered from the example in this literature, where hazards or efficacy are affected by such alterations.

EPA Est. 2724-TX-1

Manufactured by THURON INDUSTRIES, INC.
12200 Denton Drive – Dallas, Texas 75234

NET CONTENTS

LOT NO.

Package before date:

LIN. FT.

SAMPLE LABEL

DIRECTIONS FOR USE: Buckle collar around dog's neck. This collar, like all collars, should be worn loosely to prevent irritation. When collar is first placed on dog, adult ticks will be killed in a few days and will fall off or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages, ticks are smaller than a match head and are difficult to see. The Thuron Tick Collar will also control these immature stages. If ticks are a problem, this collar should be worn constantly. **Replace when effectiveness diminishes.**

Fleas — This collar is also highly effective in preventing and killing fleas. As long as your dog is wearing a Thuron Tick Collar, do not use a separate flea collar. **Replace when effectiveness diminishes.**

ACTIVE INGREDIENT:

o-Isopropoxyphenyl methylcarbamate* 9.4%

INERT INGREDIENTS: 90.6%

CAUTION: Do not open protective pouch until ready to use. Do not allow children to handle this collar. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction.* In case of contact, flush eyes with water. Wash hands thoroughly with soap and water after handling collar. The dust released by this collar is a cholinesterase inhibitor.

Note to Physician: Atropine and *homatropine are antidotal.

When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Do not use on sick or convalescing dogs. Do not use other pesticides on dog while collar is worn. Remove collar when bathing dog.