

THURON

TICK KILL COLLAR FOR DOGS

For repackaging of an insecticide only — Not for resale

See attached sample label

Confidential data to support the label are on file with the Environmental Protection Agency in file EPA Reg. No. 2724-252.

ACTIVE INGREDIENT: *o*-Isopropoxyphenyl methylcarbamate 9.4%
INERT INGREDIENTS: 90.6%

EPA REG. NO. 2724-256

CAUTION: DO NOT ALLOW CHILDREN TO HANDLE COLLAR

Dust will form on collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction. In case of contact, flush eyes with water. Wash hands thoroughly with soap and water after handling collar. The dust released by the collar is a cholinesterase inhibitor.

Note to Physician: Atropine and homatropine are antidotal.

NOTE: Additional data may be required to support registration of your product, if your labeling text or graphics are altered from the example in this literature, where hazards or efficiency are affected by such alterations.


EPA Est. 2724-TX-1

Manufactured by THURON INDUSTRIES, INC.
12200 Denton Drive - Dallas, Texas 75234

CONTENTS

LOT NO.

Package before date:



SAMPLE LABEL

DIRECTIONS FOR USE: Place collar around dog's neck. This collar, like all collars, should be worn only to prevent irritation. When collar is first put on dog, adult dogs will be cut during a few days and will fall off or may tear and be removed. Puppies can be dog for three weeks. In the first three weeks, they are smaller than a match head and are difficult to see. The Intron Tick Collar will also control these immature stages. If ticks are a problem, this collar should be worn constantly. Replace when effectiveness diminishes.

Fleas: This collar is also highly effective in preventing and killing fleas. As long as your dog is wearing a Thorex Tick Collar, do not use a separate flea collar. Replace when effectiveness diminishes.

ACTIVE INGREDIENT:

o-Propoxyphenyl methyl isothiocyanate 9.4

INERT INGREDIENTS: 90.6

CAUTION: Do not open protective pouch until ready to use. Do not allow children to handle the collar. Dust will form on this collar during storage. Do not get dust or collar in mouth, painful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction. In case of contact, flush eyes with water. Wash hands thoroughly with soap and water after handling product. The dust is irritating to the mucous membranes of the eye.

Active Ingredient: *o*-Propoxyphenyl methyl isothiocyanate

When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or a severe reaction. Collar is intended for use only as an insecticide. It is not to be taken internally by man or animal. Do not use on sick or injured animals. Do not use on other parts of body except while animal is worn. Remove collar when bandaged.