

2724-273 PM12
1071

[] Indicates optional statements

FLEA COLLAR RF-75 FOR DOGS

Contains Sevin*

Kills fleas up to 4 months

Kills ticks up to 1 month, continues to aid in tick control for an additional 1 month.

[One-piece buckle design]

[Integrated plastic buckle]

[Unique buckle design]

[New integrated one-piece buckle]

[One-piece buckle]

KEEP OUT OF REACH OF CHILDREN

CAUTION

See cautions on back

NET CONTENTS:

ACCEPTED
MAR 0 3 1985
Unit
Patented in U.S. and other countries
Use on animals for the purpose
of flea and tick control
EPA Reg. No. 2724-273

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not open inner envelope until ready to use.

Remove the collar from package and place around dog's neck, adjust for proper fit and buckle in place. [Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Since the plastic buckle is molded into the collar, the probability of neck irritation and buckle failure is reduced.] Collar must be worn loosely, but securely enough to prevent easy removal and loss of the collar. Cut off any excess length of collar, leaving ample length for expansion as the dog grows. Dispose of excess collar by discarding in trash.

Check and adjust collar periodically to assure a proper fit. [To release the collar, squeeze buckle between thumb and finger and pull sharply.]

RESULTS EXPECTED: The flea collar starts killing fleas when it is placed around the dog's neck. The fleas on the dog will be killed and new ones which may temporarily appear on the dog will also be killed during the 4 months (17 weeks) collar is worn. This product has been shown to provide effective tick control up to 1 month, and continue to aid in tick control for an additional 1 month. For continuous protection, replace collar when effectiveness diminishes.

STORAGE & DISPOSAL:

Store in original unopened container; away from children. Do not reuse container or used collar. Wrap and put in trash.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)..... 16.0%

INERT INGREDIENTS..... 84.0%

CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

Do not use on puppies under 6 weeks of age. Do not use on pregnant dogs.

Some animals may be sensitive to this collar. Any collar, when fastened too tightly, may cause skin irritation. If irritation persists after proper adjustment, remove the collar. Do not use any other pesticide on dog while collar is worn. Do not use on sick or convalescing animals.

ZOECON INDUSTRIES

A Division of Zoecon Corporation
12200 Denton Drive, Dallas, Texas 75234

*Reg. Trademark of Union Carbide Corporation for the active ingredient carbaryl.

EPA Reg. No. 2724-273
U.S. Pat. No. 3,852,416
285-C:6941A:0144A

EPA Est. 2724-TX-1
Made in USA

2724-273 PM 12
10/1

[] Indicates optional statements

FLEA COLLAR RF-75 FOR DOGS

Contains Sevin*

Kills fleas up to 4 months

Kills ticks up to 1 month, continues to aid in tick control for an additional 1 month.

[One-piece buckle design]

[Integrated plastic buckle]

[Unique buckle design]

[New integrated one-piece buckle]

[One-piece buckle]

KEEP OUT OF REACH OF CHILDREN

CAUTION

See cautions on back

NET CONTENTS:

ACCEPTED
MAR 0 8 1985
2724-273

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not open inner envelope until ready to use.

Remove the collar from package and place around dog's neck, adjust for proper fit and buckle in place. [Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Since the plastic buckle is molded into the collar, the probability of neck irritation and buckle failure is reduced.] Collar must be worn loosely, but securely enough to prevent easy removal and loss of the collar. Cut off any excess length of collar, leaving ample length for expansion as the dog grows. Dispose of excess collar by discarding in trash.

Check and adjust collar periodically to assure a proper fit. [To release the collar, squeeze buckle between thumb and finger and pull sharply.]

RESULTS EXPECTED: The flea collar starts killing fleas when it is placed around the dog's neck. The fleas on the dog will be killed and new ones which may temporarily appear on the dog will also be killed during the 4 months (17 weeks) collar is worn. This product has been shown to provide effective tick control up to 1 month, and continue to aid in tick control for an additional 1 month. For continuous protection, replace collar when effectiveness diminishes.

STORAGE & DISPOSAL:

Store in original unopened container; away from children. Do not reuse container or used collar. Wrap and put in trash.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)..... 16.0%
INERT INGREDIENTS..... 84.0%

CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

Do not use on puppies under 6 weeks of age. Do not use on pregnant dogs.

Some animals may be sensitive to this collar. Any collar, when fastened too tightly, may cause skin irritation. If irritation persists after proper adjustment, remove the collar. Do not use any other pesticide on dog while collar is worn. Do not use on sick or convalescing animals.

ZOECON INDUSTRIES
A Division of Zoecon Corporation
12200 Denton Drive, Dallas, Texas 75234

*Reg. Trademark of Union Carbide Corporation for the active ingredient carbaryl.

EPA Reg. No. 2724-273
U.S. Pat. No. 3,852,416
285-C:6941A:0144A

EPA Est. 2724-1X-1
Made in USA