

ACCEPTED

JUN 17 1964

Propoxur Flea Collar RF-101 for Cats  
EPA Reg. No. 2724-275

2724-275

Front Panel

(Optional)

Use not to  
exceed 6 months  
from acceptance.

**New** [redacted] Stretch Band

**New** [redacted] Feature

[redacted] Break-away Collar

[redacted] Expandable Collar

Unique [redacted] Feature

One-piece Buckle Design

Integrated Plastic Buckle

Unique Buckle Design

**New** Integrated One-Piece  
Buckle

[redacted] One-Piece  
Buckle

#### Alternate Directions for Use

Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Collar must be worn loosely, but securely enough to prevent easy removal. Cut off any excess length and discard in trash.

The expandable section molded into the collar will stretch under the weight of an average size cat and allow the cat to free itself if it becomes hooked or entangled by the collar. Since both the expandable section and plastic buckle are molded into the collar, the probability of neck irritation and buckle failure is reduced.

To release the collar, squeeze buckle between thumb and finger and pull sharply.

Do not use on kittens under 6 weeks of age.

EPA/RD  
RECEIVED  
JUN 16 1964

**REFUND IF NOT SATISFIED**

**DIRECTIONS:** Remove collar from package and place around cat's neck. Adjust for proper fit and buckle in place. Collar must be worn loosely but securely enough to prevent easy removal and loss. Cut off any excess length of collar, leaving ample length for expansion as the cat grows, and dispose of by discarding in trash. Check and adjust collar periodically to assure a proper fit.

Do not use on kittens under 6 weeks of age.

**RESULTS EXPECTED:** The flea and tick collar starts killing fleas and ticks when it is placed around the cat's neck. Fleas will be killed and new ones which may temporarily appear on the cat will also be killed during the 5 months. Ticks are tough occasional pests of cats and will be killed within a few days. For continuous protection, replace collar when effectiveness diminishes.

**ACTIVE INGREDIENT**

*o*-Isopropoxyphenyl methylcarbamate † ..... 9.4%

**INERT INGREDIENTS** ..... 90.6%

**CAUTION:** Do not open inner envelope until ready to use. Do not allow children to play with this collar. Dust will form on this collar during storage. Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction. In case of contact, flush eyes with water. Wash hands thoroughly with soap and water after handling collar. The dust released by this collar is a cholinesterase inhibitor.

Note to Physician: Atropine and homatropine are antidotal.

When collar is first worn, observe neck area every few days for irritation. Any collar, when fastened too tightly, may cause skin irritation. Remove collar at the first sign of irritation or adverse reaction. Collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Do not use on sick or convalescing animals. Do not use any other pesticide on cat while collar is worn.

EPA Reg. No. 2724-275

Zoecon Industries

12200 Denton Drive Dallas, Texas 75234

Made in USA

EPA Est 2724-TX-1

U.S. Patent No. 3,852,612; 3,111,539

# Propoxur Flea Collar For Cats RF-101

## Really kills fleas and ticks up to 5 months

**FOR CATS OF ALL SIZES**

NET WT. 0.29 ounces 8.35 grams

**CAUTION:** Do not allow children to play with this collar.  
See cautions on back

Front Panel

(OPTIONAL)

FREE NITE TIME SAFETY REFLECTORS See Side(s)

Left Side Panel

(OPTIONAL)

TWO FREE SAFETY REFLECTOR SLEEVES IN BOX. ATTACH REFLECTOR SLEEVES TO COLLAR AS FOLLOWS:

- (1) Clip end of collar to a point (Illustration A)
- (2) Push collar thru reflector sleeve as shown. Position silver reflector on same side of collar as buckle. (Illustration B)
- (3) Position reflector sleeves on collar as shown. (Illustration C)

